

UNIVERSIDADE DE SÃO PAULO

**INSTITUTO DE FÍSICA
CAIXA POSTAL 66318
05389-970 SÃO PAULO - SP
BRASIL**

PUBLICAÇÕES

IFUSP/P-1221

**COLOR OCTET WAVE FUNCTIONS IN HEAVY
QUARKONIA**

F.S. Navarra and C.A.A. Nunes

Instituto de Física, Universidade de São Paulo

P. Ring

Physik-Department der Technischen Universität München
D-8046 Garching, Germany

M. Schaden

Department of Physics, New York University, New York,
NY 10003, USA

Maio/

Color Octet Wave Functions in Heavy Quarkonia

F. S. Navarra and C. A. A. Nunes

Instituto de Física, Universidade de São Paulo
Caixa Postal 66318, 05389-970, São Paulo, Brasil

P. Ring

Physik-Department der Technischen Universität München
D-8046 Garching, Germany

M. Schaden

Department of Physics, New York University, New York, NY 10003, USA

Abstract

An effective hamiltonian for heavy quarkonia is derived from QCD by separating gluonic fields in background and quantum fields and neglecting anharmonic contributions. Mesonic states with nonperturbative gluonic components are constructed. These states are invariant under gauge changes of the background fields and form an orthogonal basis. The effective hamiltonian is diagonalized in this basis in a systematic $1/m$ - and short distance expansion. For very heavy quarkonia, we obtain an effective potential similar to the phenomenological funnel potential. The results to order $1/m$ for pseudoscalar meson masses and wave functions are shown and compared with those of the Cornell model. Special attention is given to the color octet wave functions which, in our approach, can be computed and not just parametrized as it is the case in most of the literature.

1 Introduction

Calculations of heavy quarkonium annihilation rates have acquired a rigorous theoretical treatment through general factorization formulae[1] obtained in the context of the effective field theory, NRQCD[2]. In the factorization formulae short-distance coefficients, calculated in perturbation theory, are combined with long-distance matrix elements that can so far only be computed in lattice simulations. In spite of its limitations, this approach provided the solution of the long-standing problem of infrared divergences in h_c and χ_{c1} decays into light hadrons. The probability for the $q\bar{q}$ pair to be in a color-octet S-wave at the origin is an essential input in this approach. Since accurate lattice calculations of this probability are not yet available, it is determined phenomenologically[3]. NRQCD has also given a good explanation for the production rate of prompt J/Ψ and Ψ' at large transverse momentum at the Tevatron[4], which is larger than previous theoretical expectations. At such energies fragmentation of partons into quarkonium states becomes an important component of quarkonium production[5]. Fragmentation of a gluon into a $c\bar{c}$ pair in a pointlike color-octet S-wave state of χ_{cJ} mesons with a cascade decay to J/Ψ is crucial to bring theoretical predictions for J/Ψ production closer to experimental data[6]. It has been suggested[7] that the puzzling Ψ' surplus (a factor of 30) could be explained by the fragmentation of a gluon into a pointlike color-octet S-wave state. In any case, color-octet matrix elements are needed. In the framework of NRQCD, since one can not compute these matrix elements, they have to be first extracted from some experiments and then used to study some other processes.

In this work, in the same way as in NRQCD, we deal with octet states of the type $|q\bar{q}g\rangle$, where g is a soft gluon, which is, in a first approximation, indistinguishable from the non-trivial vacuum gluons. In the approach we propose, the probability for the $q\bar{q}$ pair to be in a color octet state can in principle be computed, since such octet configurations are included in the description of the meson from the outset. In this sense, our approach could be considered an inexpensive alternative to lattice simulations.

We calculate the short distance expansion of the hamiltonian matrix elements in a gauge invariant basis of color singlet states[8], including only terms up to order r^2 and $1/m$. We then obtain the eigenstates by (numerical) diagonalization of this hamiltonian matrix and see how far we can go in this approach. The previous discussion implies that a realistic description of

heavy quarkonia within this short distance expansion can only be obtained if the (background) gauge invariant basis of ref.[8] is extended to include states in which the heavy quark-antiquark pair is in a color octet representation but coupled to a nonperturbative gluonic background field[9, 10] to form an overall color singlet. Diagonalization of the hamiltonian matrix leads to a “color-octet” component in the wavefunction of heavy quarkonia that takes into account the possibility of color exchange between the valence quarks and the background field. This component becomes rather large for higher excited states and its coupling to the “singlet” is the main reason for the distortion of the coulombic spectrum in this model. In the pseudoscalar meson channel only the few octet states constructed in section 2 couple and a numerical diagonalization of the resulting system of differential equations is still quite feasible.

In section 3 we derive the effective hamiltonian appropriate for the description of heavy quarkonia. We first obtain the effective Lagrangian to order $1/m$ by a Foldy-Wouthuysen[11] transformation. Using the background field formalism[12] and neglecting anharmonic quantum fluctuations one finally arrives at an effective Lagrangian[13, 14] that includes background fields up to 2nd order and which, in the instantaneous approximation, gives rise to an effective hamiltonian that is accurate to order $1/m$ and r^2 . Since retardation effects are of order $1/m^2$, their consistent inclusion would require a much more elaborated treatment, which would only obscure the basic non-perturbative gluonic effects that we want to elucidate here.

Numerical diagonalization of this effective hamiltonian in the extended basis then yields (pseudoscalar) meson masses and wave functions. Our approach can in principle not be described by an effective potential in the singlet channel[9, 10] because the elimination of the relative octet states would make it energy dependent. One can however obtain an energy independent effective potential for infinitely heavy quarks, where all terms of order $1/m$ (also the kinetic energy) can be neglected. This potential should be closely related to the static quark potential that one extracts from the Wilson loop[15]. This potential, which we derive and discuss in section 4, also gives us an idea of how far the short distance expansion can be trusted. We show that it is apparently linearly rising at intermediate distances $.4fm < r < .7fm$.

In section 5 we compare our method to methods relying on 2nd-order perturbation theory in the background field and to the phenomenological Cornell[16] potential. We first present results where all potential matrix

elements of order $1/m$ are neglected. This greatly simplifies the calculations because only a few basis states couple, but pseudoscalar and vector mesons are degenerate at this level. Our formalism with this approximation has been recently applied to the study of temperature effects on charmonium masses [17]. We end the section presenting results of a more complete calculation of the pseudoscalar quarkonia which includes the $1/m$ corrections.

Section 6 is a summary and discussion of our results.

2 Gauge Invariance and Basis States

It is usually assumed that physical states are color singlets. For heavy mesons, where the non-relativistic approximation is adequate, one can represent quark and antiquark fields by 2-component spinors. Gauge invariance of the state requires that color is parallel transported from the quark to the anti-quark along some path. With a straight path the quark and anti-quark anti-commutation relations ensure that the basis states are orthogonal[8]. For simplicity we will restrict ourselves in the following to pseudoscalar mesons. The simplest gauge invariant pseudoscalar state is of the form

$$|21\rangle_S = \frac{1}{\sqrt{6}} \sum_{ab,\alpha} u_\alpha^{\dagger a}(\vec{x}_2) T_{ab}(\vec{x}_2, \vec{x}_1) v_\alpha^b(\vec{x}_1) |\Omega\rangle, \quad (1)$$

where $u^{\dagger a}(\vec{x}_2)$ creates a quark with color a and spin α at \vec{x}_2 . v does the same for an antiquark. We will refer to such gauge invariant states as singlet states since the quark anti-quark pair at vanishing separation is in a color singlet representation.

We choose for the color transport operator

$$T_{ab}(\vec{x}_2, \vec{x}_1) = Pexp(-ig \int_{\vec{x}_1, t}^{\vec{x}_2, t} dx^\mu A_\mu(x))_{ab}, \quad (2)$$

the path ordered exponential (denoted by $Pexp$) of gluon operators $A_\mu(x)$ along a straight line from \vec{x}_1 to \vec{x}_2 . It is then relatively straightforward to show that canonical anti-commutation relations for the quark and anti-quark operators imply that the singlet meson states (7) are orthogonal[8].

We can also construct gauge invariant basis states where the valence quark anti-quark pair is in a color octet representation at vanishing separation by coupling them to chromo-electric or -magnetic fields. We shall

call these states octet states for obvious reasons. Since the chromomagnetic field $\vec{B} = \vec{B}^a \lambda^a / 2$ transforms as a pseudovector and the chromoelectric field $\vec{E} = \vec{E}^a \lambda^a / 2$ as a vector under rotations and according to the adjoint representation of the gauge group, we extend the basis for pseudoscalar mesons by the states

$$|21\rangle_B = \sum_{\alpha\beta} \frac{g}{\pi\phi} u_\alpha^\dagger(\vec{x}_2) \vec{\sigma}_{\alpha\beta} \cdot \vec{B}(\vec{x}_2, t) T(\vec{x}_2, \vec{x}_1) v_\beta(\vec{x}_1) |\Omega\rangle, \quad (3)$$

$$|21\rangle_{E1} = \sum_{\alpha} \frac{\sqrt{3}g}{\pi\phi} u_\alpha^\dagger(\vec{x}_2) \vec{E}(\vec{x}_2, t) \cdot (\vec{x}_2 - \vec{x}_1) T(\vec{x}_2, \vec{x}_1) v_\alpha(\vec{x}_1) |\Omega\rangle, \quad (4)$$

and

$$|21\rangle_{E2} = \sum_{\alpha\beta} \frac{i\sqrt{3}g}{\sqrt{2}\pi\phi} u_\alpha^\dagger(\vec{x}_2) \vec{E}(\vec{x}_2, t) \cdot (\vec{\sigma}_{\alpha\beta} \times (\vec{x}_2 - \vec{x}_1)) T(\vec{x}_2, \vec{x}_1) v_\beta(\vec{x}_1) |\Omega\rangle, \quad (5)$$

the summation over color indices being implied. The gluon fields in eqs. (3)-(5) are fields of soft gluons. High momentum gluons are expected not to be important for the low-lying states. The separation in soft (or background) and high (or quantum) momentum fields is made explicit in the next section.

The above states are seen to be mutually orthogonal and normalized by the canonical anti-commutation relations of the quark and anti-quark operators if we assume expectation values

$$\begin{aligned} \langle \Omega | \frac{g^2}{4\pi^2} B^{ia} B^{jb} | \Omega \rangle &= -\langle \Omega | \frac{g^2}{4\pi^2} E^{ia} E^{jb} | \Omega \rangle \\ &= \frac{1}{96} \delta^{ij} \delta^{ab} \langle \Omega | \frac{\alpha}{\pi} F^{\mu\nu c} F_{\mu\nu}^c | \Omega \rangle = \frac{1}{96} \delta^{ij} \delta^{ab} \phi^2, \end{aligned} \quad (6)$$

and $\langle E \rangle = \langle B \rangle = \langle EB \rangle = 0$, which is a consequence of the Lorentz- and parity- invariance of the vacuum state $|\Omega\rangle$. Its nontrivial nature is reflected in a non-vanishing value for ϕ^2 , which from QCD sumrule estimates should be close to $(330 \text{ MeV})^4$ [18]. If we neglect expectation values of all higher dimensional gluonic operators the states (1),(3),(4) and (5) form a complete orthogonal basis for the valence quarks of heavy pseudoscalar mesons, while the singlet states (1) are only complete in this sense if one neglects the non-trivial vacuum structure altogether.

We thus effectively truncate the Fock space by taking basis states whose gluonic sector contains at most 1 soft gluonic mode, i.e. we replace the low momentum gluonic modes with octet quantum numbers by zero-modes. We show in section 3 that such a severely truncated basis is adequate for the problem at hand. Lorentz invariance of the ground state $|\Omega\rangle$ forces one to assume that the chromoelectric background field is either antihermitian or that the “E-states” have negative norm (see (6)). Both possibilities lead to a nonhermitian hamiltonian matrix, whose eigenvalues in general are not real. Already the construction of basis states for the mesons on a non-trivial ground state indicates that one can at best hope to find a few stable mesons in this approach. We assume that a pseudoscalar quarkonium η is well described by a linear combination of the above basis states

$$|\eta\rangle = \sum_{M=S,E1,E2,B} \int_{1,2} \psi_M(2,1) |2,1\rangle_M. \quad (7)$$

3 Effective Hamiltonian

We would like to use the basis constructed in section 2 to approximately diagonalize the QCD hamiltonian for heavy quarkonia. If low lying heavy quarkonia were of sufficient small size, the interaction would be dominated by the perturbative gluon exchange, which would lead to a Coulomb-like effective potential. The experimental spectrum deviates however noticeably from a purely coulombic one and can be reasonably well reproduced by the combination of a Coulomb with a linear confining force. We wish to emphasize here that the truly long range part of the confining force ($r > 1\text{fm}$) is not really tested by the observed quarkonium states [19]. Our conjecture is, that what has to be included in a systematic approach are the first short distance corrections to the (perturbative) Coulomb force due to the non-trivial structure of the gluonic ground state. This was proposed previously but without any tangible results, because the corrections were found to be exceedingly large within the framework of 2nd-order perturbation theory except for quarkonia beyond bottomium [9, 20]. This apparent failure of an idea which we believe is quite well founded, has led us to reexamine the basic procedure used in the evaluation of these effects.

In this section we outline the derivation of the effective hamiltonian for heavy quarkonia, whose matrix elements are correct up to order $\alpha, 1/m$ and

r^2 . To this order in the short distance and heavy quark expansion the non-perturbative aspects of the ground state can be described by the gluon condensate ϕ^2 .

The nonrelativistic approximation for heavy quarks is conveniently obtained from the QCD lagrangian

$$\mathcal{L}(x) = -\frac{1}{4}F_{\mu\nu}^a F_a^{\mu\nu} + \bar{\psi}(i\cancel{D} + gT^a V_a)\psi - m\bar{\psi}\psi, \quad (8)$$

by a Foldy-Wouthuysen transformation[11]. In terms of transformed quark fields

$$\begin{aligned} \psi &\rightarrow \exp(i\vec{\gamma} \cdot \vec{D}/2m)\psi, \\ \bar{\psi} &\rightarrow \bar{\psi} \exp(-i\vec{\gamma} \cdot \vec{D}/2m), \end{aligned} \quad (9)$$

the lagrangian is

$$\begin{aligned} \mathcal{L}_{NRQCD}(x) &= -\frac{1}{4}F_{\mu\nu}^a F_a^{\mu\nu} + \bar{\psi}(i\gamma^0 D_0 - m)\psi \\ &\quad + \bar{\psi}\left(\frac{\gamma^0}{2m}[\vec{\gamma} \cdot \vec{D}, D_0] + \frac{(i\vec{\gamma} \cdot \vec{D})^2}{2m}\right)\psi + O(1/m^2) \\ &= -\frac{1}{4}F_{\mu\nu}^a F_a^{\mu\nu} + \bar{\psi}(i\gamma^0 D_0 - m)\psi + \bar{\psi}\frac{\vec{D}^2}{2m}\psi \\ &\quad - \bar{\psi}\frac{ig\vec{\alpha} \cdot \vec{E}}{2m}\psi + \bar{\psi}\frac{g\vec{\Sigma} \cdot \vec{B}}{2m}\psi + O(1/m^2), \end{aligned} \quad (10)$$

where $\vec{D} = \vec{\partial} - ig\vec{V}$ and $\vec{\Sigma} = \begin{pmatrix} \vec{\sigma} & 0 \\ 0 & \vec{\sigma} \end{pmatrix}$ does not couple upper and lower spinor components. They are only coupled in order $1/m$ by the $\vec{\alpha}$ matrices (which appear in the usual Dirac Hamiltonian). The above lagrangian can be further reduced to order $1/m^2$ by another transformation

$$\psi \rightarrow \exp(-ig\vec{\alpha} \cdot \vec{E}/4m^2)\psi, \text{ etc.} \quad (11)$$

The non-relativistic lagrangian to order $1/m$ finally is

$$\mathcal{L}_{NRQCD}(x) = -\frac{1}{4}F_{\mu\nu}^a F_a^{\mu\nu} + \bar{\psi}(i\gamma^0 D_0 - m)\psi + \bar{\psi}\frac{\vec{D}^2}{2m}\psi + \bar{\psi}\frac{g\vec{\Sigma} \cdot \vec{B}}{2m}\psi, \quad (12)$$

which can be written in terms of uncoupled 2-component spinors by decomposing $\psi = (u, v^\dagger)$ $\bar{\psi} = (u^\dagger, -v)$ and using the Dirac representation of the γ -matrices.

In order to perform a short distance expansion we separate the gluonic fields in slowly varying background-[12](A) and quantum- (Q) fields having high Fourier components:

$$V_\nu^a = A_\nu^a + Q_\nu^a. \quad (13)$$

The division (13) can however only be defined if the gauge is fixed. In deriving the spin-dependence of nonperturbative interactions for heavy quarkonia Curci et al.[13] found a particular gauge very convenient. We essentially follow their procedure here and impose the Coulomb background gauge condition

$$D_i Q^i = 0 \quad (14)$$

for the quantum fields, where

$$D_\mu Q_\nu = \partial_\mu Q_\nu + gA_\mu \times Q_\nu = \partial_\mu Q_\nu + gf^{abc}A_{\mu b}Q_{\nu c}. \quad (15)$$

The background fields are defined in a modified Schwinger gauge[13]

$$A_j^b = -\frac{1}{2}F_{ji}^b x^i \quad ; \quad A_0^b = -F_{0i}^b x^i, \quad (16)$$

valid to order x^2 , where we assumed that the field-strengths corresponding to the background fields are constant (or have sufficiently low momenta, such that they can be regarded as essentially constant over the extent of the meson). This definition of the background fields in terms of (practically) constant field strengths also gives a precise meaning to the separation in equation (13). It also implies that the background fields in this gauge are (practically) time independent - a property which will become useful when a Hamiltonian is required.

We next expand the nonrelativistic Lagrangian only to second order in the quantum fields and subsequently integrate them out in favor of an effective (coulombic) interaction. Although ghost terms are necessary in the gauge defined by (14), they do not contribute to quadratic order in the quantum fields. Our truncation of the interaction terms for the quantum fields eliminates all radiative corrections. To obtain them one would have to go beyond this approximation and calculate perturbative corrections before eliminating hard gluons. Fortunately, the asymptotic freedom of QCD guarantees that they are only logarithmic at short distances and could be accounted for by a

running coupling constant. These logarithmic corrections to the Coulomb potential are not dramatically important for describing heavy quarkonia spectra. We show their effect in section 5. The truncation of the interaction terms for the radiative corrections eliminates also n-body potential terms which could be of importance for the description of baryons.

The matrix elements of the slowly varying background fields (A) will however be parametrized. Their amplitude is large and an expansion in the coupling in this case not applicable.

After the elimination of the quantum gluonic fields by their equations of motion the effective lagrangian in terms of the background and heavy quark fields becomes[14]:

$$\begin{aligned}
L_{eff} = & \int d^4x \left\{ -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \bar{\psi} (i\gamma^0 \partial_0 + g\gamma^0 A_0 - m) \psi \right. \\
& + \frac{1}{2m} \bar{\psi} (\vec{\nabla}^2 - ig\vec{\nabla} \cdot \vec{A} - ig\vec{A} \cdot \vec{\nabla} - g^2 \vec{A}^2) \psi + \frac{g}{2m} \bar{\psi} \vec{\Sigma} \cdot \vec{B} \psi \\
& + g^2 \int d^4y \psi^\dagger(x) T^a \psi(x) \mathcal{D}^{ab}(x, y) \psi^\dagger(y) T^b \psi(y) \\
& \left. - \frac{1}{2} \int d^4y J_i^a(x) (\tilde{\mathcal{D}}_{ij}^{ab}(x, y) - \mathcal{K}_{ij}^{ab}(x, y)) J_j^b(y), \right. \quad (17)
\end{aligned}$$

where

$$\begin{aligned}
J_k^a(x) = & -\frac{ig}{2m} [(\partial_k \bar{\psi}) T^a \psi - \bar{\psi} T^a \partial_k \psi - i\epsilon_{ijk} \partial_i (\bar{\psi} T^a \Sigma_j \psi)] \\
& + \frac{g^2}{2m} \bar{\psi} (T^b T^a + T^a T^b) \psi A_k^a - \frac{g^2}{2m} f^{abc} \epsilon_{ijk} A_i^b \bar{\psi} T^c \Sigma_j \psi \\
& - 2g^2 f^{adc} F_{k0}^d \int d^4z \mathcal{D}^{cb}(x, z) \psi^\dagger(z) T^b \psi(z). \quad (18)
\end{aligned}$$

The propagator \mathcal{D} relates the Q_0 field to its source

$$Q_0^a(x) = \int d^4y \mathcal{D}^{ab}(x, y) j_b^0(y), \quad (19)$$

with

$$j_0^a = g\psi^\dagger T^a \psi + 2gf^{abc} F_{b0}^c Q_{c0}, \quad (20)$$

and satisfies

$$(D_j D_j)^{ab} \mathcal{D}^{bd}(x, y) = \delta^{ad} \delta^4(x - y). \quad (21)$$

Similarly the propagator $\tilde{\mathcal{D}}$ appears in the elimination of the spatial components Q_i and satisfies

$$\int d^4z M_{ij}^{ab}(x, z) \tilde{\mathcal{D}}_{jk}^{bc}(z, y) = \delta^{ac} \delta^4(x - y) \delta_{ik}, \quad (22)$$

with the differential operator M_{ij}^{ab} given by

$$\begin{aligned}
M_{ij}^{ab}(x, z) = & \delta^4(x - z) [-(D_\mu D_\mu)^{ab} \delta_{ij} - 2gf^{abc} F_{ij}^c \\
& - \frac{g^2}{2m} \bar{\psi} (T^a T^b + T^b T^a) \psi \delta_{ij} + \frac{g^2}{2m} f^{abc} \epsilon_{ijk} \bar{\psi} T^c \Sigma^k \psi] \\
& + 4g^2 f^{da} f^{cb} F_{i0}^d(x) \mathcal{D}^{dc}(x, z) F_{j0}^e(z). \quad (23)
\end{aligned}$$

Finally, the propagator \mathcal{K} enters when the Lagrange multiplier of the gauge fixing condition for the quantum fields (14) is eliminated in turn. Its equation of motion is

$$\int d^4z D_i^{ba}(x) \tilde{\mathcal{D}}_{ij}^{ad}(x, z) D_j^{dc}(z) \mathcal{K}^{cd}(z, y) = \delta^{bd} \delta(x - y). \quad (25)$$

These rather formidable integro-differential equations for the Green functions can formally be solved order by order in the background field A . Since we will only retain terms of the hamiltonian matrix proportional to the lowest dimensional condensate $\langle g^2 FF \rangle$, we only keep terms up to second order in the background fields in the gauge (16). As we will see, linear terms in the background field have to be retained, although we will assume that $\langle F \rangle = 0$, e.g. that global colour- and Lorentz- invariance is not broken.

In order to obtain a tractable hamiltonian, further approximations are necessary. We will neglect all retardation effects in the effective interaction. This instantaneous approximation is correct to order $1/m$, because retardation effects are generally expected to be of order $1/m^2$. Since the interaction is instantaneous, the effective hamiltonian becomes time independent even in the presence of the background fields (which are (nearly) time independent in the gauge (16)). This greatly simplifies the interpretation of our results.

The effective hamiltonian, correct to order $1/m$, r^2 and α_s therefore is

$$\begin{aligned}
H = & \int d^3x \left\{ u^\dagger(\vec{x}) m u(\vec{x}) + v(\vec{x}) m v^\dagger(\vec{x}) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right. \\
& \left. - u^\dagger(\vec{x}) T^A g E_i^A x_i u(\vec{x}) - v(\vec{x}) \bar{T}^A g E_i^A x_i v^\dagger(\vec{x}) \right.
\end{aligned}$$

$$\begin{aligned}
& -u^\dagger(\vec{x})\frac{\vec{\nabla}_{\vec{x}}^2}{2m}u(\vec{x}) - v(\vec{x})\frac{\vec{\nabla}_{\vec{x}}^2}{2m}v^\dagger(\vec{x}) \\
& +\alpha\int d^3y u^\dagger(\vec{x})T^A u(\vec{x})\frac{1}{r}v(\vec{y})\bar{T}^A v^\dagger(\vec{y}) \\
& +\frac{\alpha}{2}\epsilon_{ijk}gB_k^C f^{ACB}\int d^3y u^\dagger(\vec{x})T^A u(\vec{x})\frac{y_j x_i}{r}v(\vec{y})\bar{T}^A v^\dagger(\vec{y}) \\
& -\frac{\pi^2\phi^2\alpha}{64}\int d^3y u^\dagger(\vec{x})T^A u(\vec{x})\frac{(\vec{x}\times\vec{y})^2}{r}v(\vec{y})\bar{T}^A v^\dagger(\vec{y}) \\
& +\frac{1}{m}\left[\frac{\pi^2\phi^2}{128}u^\dagger(\vec{x})u(\vec{x})\vec{x}^2 + \frac{\pi^2\phi^2}{128}v(\vec{x})v^\dagger(\vec{x})\vec{x}^2\right. \\
& -\frac{i}{2}u^\dagger(\vec{x})T^A\epsilon_{ijk}gB_k^A x_j\partial_{x_i}u(\vec{x}) - \frac{i}{2}v(\vec{x})\bar{T}^A\epsilon_{ijk}gB_k^A x_j\partial_{x_i}v^\dagger(\vec{x}) \\
& -\frac{1}{2}u^\dagger(\vec{x})\sigma_i T^A gB_i^A u(\vec{x}) - \frac{1}{2}v(\vec{x})\sigma_i \bar{T}^A gB_i^A v^\dagger(\vec{x}) \\
& +\frac{i\alpha}{8}f^{ADC}gE_j^D\left[(\partial_{x_i}u^\dagger(\vec{x}))T^A u(\vec{x}) - u^\dagger(\vec{x})T^A(\partial_{x_i}u(\vec{x}))\right. \\
& \quad \left.- i\epsilon_{ijk}\partial_{x_i}(u^\dagger(\vec{x})T^A\sigma^k u(\vec{x}))\right]\cdot\int d^3y(-3\delta_{ij}r + \frac{r_i r_j}{r})v(\vec{y})\bar{T}^C v^\dagger(\vec{y}) \\
& -\frac{i\alpha}{8}f^{ADC}gE_j^D\left[-(\partial_{x_i}v(\vec{x}))\bar{T}^A v^\dagger(\vec{x}) + v(\vec{x})\bar{T}^A(\partial_{x_i}v^\dagger(\vec{x}))\right. \\
& \quad \left.+ i\epsilon_{ijk}\partial_{x_i}(v(\vec{x})\bar{T}^A\sigma^k v^\dagger(\vec{x}))\right]\cdot\int d^3y(-3\delta_{ij}r + \frac{r_i r_j}{r})u^\dagger(\vec{y})T^C u(\vec{y})\left.\right\}.
\end{aligned} \tag{26}$$

Here $u(\vec{x})$ and $v(\vec{x})$ denote the annihilation operators for a quark and antiquark of mass m respectively whose spin and color indices have been suppressed, $\vec{r} = \vec{x} - \vec{y}$ and T^A, \bar{T}^A are the hermitian generators of the SU(3) color Lie-algebra in the 3 and $\bar{3}$ representation.

This is essentially the Hamiltonian to order $1/m$ derived previously by Curci et al.[13] in position space, except for a term which can be regarded as a long range correction of the Coulomb potential

$$\frac{3\pi^2\phi^2\alpha}{64}\int d^3x d^3y u^\dagger(\vec{x})T^A u(\vec{x})r^3 v(\vec{y})\bar{T}^A v^\dagger(\vec{y}). \tag{27}$$

We have disregarded this term because it is of order r^3 and our gauge fixing condition (16) is not valid at this order.

The terms of the Hamiltonian (26) linear in the chromo -electric and -magnetic fields as well as those proportional to \vec{x}^2 are obviously also not

translationally invariant and therefore apparently depend on the chosen gauge fixing point in (16). This gauge dependence of (26) is however absent[14] in its matrix elements in the basis (1),(3),(4) and (5). It cancels against terms which appear when derivatives act on the color transport matrix of the states. This cancelation of course only works to a certain order in the short distance expansion and only occurs if the Hamiltonian and the basis in which it is diagonalized are defined consistently. This is not a big surprise in any gauge theory, but a gauge invariant scheme to include nonperturbative aspects of the ground state in a hamiltonian formulation without losing gauge invariance was only first proposed in [8], but not used to its full extent there. From the above we see that a short distance expansion in the construction of the basis as well as the hamiltonian to order r^2 can be performed and the effects from a nontrivial expectation value ϕ^2 consistently included.

The straightforward but lengthy computation of the matrix elements of the Hamiltonian (26) in the basis for pseudoscalar mesons (7) to obtain the coupled differential equations (30) for the wavefunctions[14] will not be exhibited here. We do however also have to account for matrix elements of the purely gluonic part of the Hamiltonian

$$H_G = \int d^3x \mathcal{H}_G(x), \quad \mathcal{H}_G(x) = (E^2 + B^2)/2 + O(\alpha) \tag{28}$$

in the basis states (1),(3),(4) and (5). The order α terms in (28) are due to the 1-loop corrections from the quantum fields. When these corrections are properly included, the above Hamiltonian should be consistent with the trace anomaly[22] and reproduce the relation between the energy density of the nonperturbative gluonic ground state and the condensate value. All we will need in what follows is that this correction is a local operator of dimension 4 and order α (such as $(11\alpha/16\pi)(E^2 - B^2)$).

From rotational symmetry we conclude that only diagonal matrix elements of H_G can be non-vanishing. Since we neglect expectation values of operators with dimension greater than 4, matrix elements in "octet"-states (3),(4),(5) vanish in this approximation.

From (6) we might naively infer that the matrix elements of H_G between "singlet" states vanish as well. Phenomenologically we do however need an energy splitting between the "singlet"- and "octet"- states. Leutwyler[23] attributes it to an effective mass of the low frequency gluonic modes. A more careful examination of contributions from H_G reveals the origin of such

an effective mass in the present approach. To obtain the energy of the meson relative to that of the vacuum, one has to commute the purely gluonic hamiltonian with the creation operator for the meson. As discussed previously, gauge invariance requires that this creation operator contains a gluonic string, or color transport matrix. Within our approximation, we therefore have to consider the commutator of the creation operator for the singlet component with the local Hamiltonian density of $\mathcal{H}_G(x)$. Defining the creation operator for the singlet component of the meson by

$$\int_{1,2} \psi_S(1,2) |2,1\rangle_S = O_S^+ | \Omega \rangle,$$

the contribution of the lowest dimensional operator to the commutator is of the form

$$[\mathcal{H}_G(x), O_S^+] = O_S^+ A(x) \theta_V(x),$$

where $A(x)$ is a gauge invariant scalar operator of dimension 4 and order α (because the g^0 -part of the string obviously commutes) and $\theta_V(x)$ is a dimensionless c-number function that vanishes for x outside the meson (because the string in O_S^+ and x are otherwise separated by a spacelike distance). Since we do not neglect the vacuum expectation of the gauge invariant operator αF^2 in our approach, the matrix element in the singlet channel within our approximation becomes

$$\int d^3x_S \langle 2,1 | [\mathcal{H}_G(x), O_S^+] | \Omega \rangle = C \Psi_S(1,2), \quad (29)$$

where C is a constant which depends on the exact nature of the function $\theta_V(x)$.

Although we cannot determine the value of this constant on theoretical grounds, the above argument shows that it would be inconsistent with our approximations to neglect this contribution of H_G in the singlet channel. A non-vanishing constant C implies that the lowest order energies of the singlet and octet states differ by gluonic contributions. Our best fit to the quarkonium spectrum was obtained with $C = -756$ MeV. $m_g = -C$ can also be interpreted as an effective mass for the low frequency gluon mode in the octet-channel (since a small change in the overall normalization of the energy can be absorbed in a slight change of the heavy quark mass). This is the point of view taken by Leutwyler[23]. Our estimate $m_g = -C \sim 756$ MeV is

not in conflict with the fact that the lightest glueball has at least two such excitations and a mass of ~ 1700 MeV[24], although a somewhat smaller glueball mass would seem more natural. States with two soft effective gluons would be much higher in energy than the first states of quarkonia and should therefore have very small amplitudes.

Including this energy shift between ‘‘singlet’’ and ‘‘octet’’ components, the coupled set of differential equations for flavor singlet pseudoscalar mesons becomes

$$\begin{aligned} \left[2m - E + C - \frac{1}{m} \frac{\partial^2}{\partial r^2} - \frac{4\alpha_s}{3r} + \frac{\pi^2 \phi^2 r^2}{36m} \right] S(r) &= \\ &+ \left(-\frac{\pi \phi r}{3\sqrt{2}} + \frac{g^2 \phi}{4\sqrt{2}m} + \frac{g^2 \phi r}{8\sqrt{2}m} \frac{\partial}{\partial r} \right) E_1(r) - \frac{5g^2 \phi}{64m} E_2(r) + \frac{\pi \phi}{\sqrt{6}} B(r) \\ \left[2m - E - \frac{1}{m} \left(\frac{\partial^2}{\partial r^2} - \frac{2}{r^2} \right) + \frac{1\alpha_s}{6r} \right] E_1(r) &= \left(\frac{\pi \phi r}{3\sqrt{2}} + \frac{g^2 \phi}{8\sqrt{2}m} - \frac{g^2 \phi r}{8\sqrt{2}m} \frac{\partial}{\partial r} \right) S(r) \\ \left[2m - E - \frac{1}{m} \left(\frac{\partial^2}{\partial r^2} - \frac{2}{r^2} \right) + \frac{1\alpha_s}{6r} \right] E_2(r) &= \frac{5g^2 \phi}{64m} S(r) \\ \left[2m - E - \frac{1}{m} \frac{\partial^2}{\partial r^2} + \frac{1\alpha_s}{6r} \right] B(r) &= \frac{\pi \phi}{\sqrt{6}} S(r). \end{aligned} \quad (30)$$

where E is the mass eigenvalue of the quarkonium and m is the mass of the constituent quarks. The functions $S(r)$, $E_1(r)$, $E_2(r)$ and $B(r)$ are related to the wave function components in the expansion (7) via

$\psi_S(r) = \frac{1}{r} S(r)$; $\psi_{E_1}(r) = \frac{1}{r^2} E_1(r)$; $\psi_{E_2}(r) = \frac{1}{r^2} E_2(r)$; $\psi_B(r) = \frac{1}{r} B(r)$. As indicated above, these equations only depend on the relative coordinate r and all reference to a gauge fixing point—present in (26)—has disappeared in the evaluation of the effective Hamiltonian in the gauge invariant basis.

Apart from the constituent quark mass, the only parameters that enter equations (30) are the (reasonably well known) condensate ϕ^2 and the constant C , whose value we have tried to estimate above, as well as the strong coupling constant α_s .

4 ‘‘Effective Potential’’

Before solving the coupled set of equations (30) by numerical methods, it is instructive to consider the limit of infinitely heavy quarkonia. In this

case, all terms proportional to $1/m$ (including the kinetic energy) in (30) can be dropped, and the resulting equations give the binding energy $V = E - 2m$ for states where the quark and anti-quark are localized a distance r apart (i.e. for wave-functions $S(r), E1(r), E2(r)$ and $B(r)$ all proportional to $\delta(r - r_0)$). It is natural to compare this mass-independent binding energy to phenomenological potentials and those extracted from the expectation values of Wilson loops in numerical lattice simulations.

In this limit the $E2$ - and B -components decouple and one has to solve the algebraic equations

$$\begin{aligned} (-V + C - \frac{4\alpha_s}{3r})S(r) &= -\frac{\pi\phi r}{3\sqrt{2}}E_1(r) \\ (-V + \frac{1}{6}\frac{\alpha_s}{r})E_1(r) &= \frac{\pi\phi r}{3\sqrt{2}}S(r). \end{aligned} \quad (31)$$

The eigenfunctions are obviously localized and the eigenvalue or effective potential, $V(r)$, given by

$$V(r) = -\frac{1}{2}\left(\frac{7\alpha_s}{6r} + \sqrt{\frac{9\alpha_s^2}{4r^2} - \frac{9\alpha_s C}{3r} - \frac{2\pi^2\phi^2 r^2}{9} + C^2}\right) \quad (32)$$

Figure 1(a) shows this effective potential and its Coulomb part for parameters $\phi^2 = (360\text{MeV})^4$, $C = -756\text{MeV}$ and $\alpha_s = 0.39$, which we found appropriate for charmonium.

Although this potential is certainly no longer valid for $r > 0.9\text{fm}$, where the root in (32) becomes purely imaginary it does show a nearly linear behaviour for intermediate distances $0.4\text{fm} < r < 0.8\text{fm}$, with a correspondingly constant force of $\sim 840\text{MeV}/\text{fm}$, which compares favorably with a string tension of about $800 - 1000\text{MeV}/\text{fm}$ extracted from recent lattice calculations of the potential[25]. In Fig. 1(b) we compare our effective potential (32) to that extracted from lattice data[25] and to the phenomenological potential used by ref[16].

It is encouraging that our rather crude approximations to the vacuum structure seem to qualitatively reproduce the potential for very heavy quarks at small distances. The analytical results of this section justify the numerical calculation of ‘‘octet’’-components in heavy quarkonia which we now present.

5 Numerical Results

Going beyond the static approximation by including the kinetic energy of the quarks but still neglecting coupling terms of order $1/m$ in (30), the quarkonium spectrum becomes discrete and the following coupled set of differential equations must be solved numerically

$$\begin{aligned} \left[2m - E + C - \frac{1}{m}\frac{\partial^2}{\partial r^2} - \frac{4\alpha_s}{3r}\right]S(r) &= -\frac{\pi\phi r}{3\sqrt{2}}E_1(r) \\ \left[2m - E - \frac{1}{m}\left(\frac{\partial^2}{\partial r^2} - \frac{2}{r^2}\right) + \frac{1}{6}\frac{\alpha_s}{r}\right]E_1(r) &= \frac{\pi\phi r}{3\sqrt{2}}S(r). \end{aligned} \quad (33)$$

This is essentially Leutwyler’s[20] approximation to the problem, who obtained the perturbation of the Coulomb spectrum in 2nd-order of ϕ and found that it is exceedingly large for canonical values of the condensate. Note that the Coulomb force is repulsive in the ‘‘octet’’ channel and 1/8-th in strength compared to the ‘‘singlet’’ channel (this is just the ratio of $T_1^a T_2^a$ in the two representations) and the $E2$ - and B -states still decouple in this approximation. In this approximation, vector- and pseudoscalar- quarkonia are furthermore still degenerate. This is expected, since the spin splitting is of order $1/m$. It is however another nontrivial consistency check of our method, because the basis states for vector mesons are of course quite different. Nevertheless equivalent equations to (33) result, if $1/m$ potential terms are neglected.

The relative minus sign of the two off-diagonal coupling terms in (33) shows that this Hamiltonian is not hermitian and that its eigenvalues will generally not be real. This effect is completely missed if the coupling is treated as a perturbation. To any finite order in perturbation theory the correction to the Coulomb spectrum is real. Perturbation theory does however show that a few (low lying) eigenvalues of (33) are real for a sufficiently small ϕ . For the canonical value $\phi^2 = (300 - 380\text{MeV})^4$ we find numerically 2 – 4 stable bound quarkonia in the pseudoscalar channel, depending on the heavy quark mass m .

But even for these low states, the deviation of the exact correction to the Coulomb eigenvalue from the 2nd-order estimate is large for the charmonium and bottomonium systems as shown in table 1. We conclude with Leutwyler[20] that perturbation theory is not applicable for canonical values of ϕ , but that

the effective non-hermitian Hamiltonian equations (33) do yield reasonable corrections in an exact solution.

The value $C = -756\text{MeV}$ used by us was adjusted to reproduce the correct splitting between η_c and η'_c [26]. The corrections to the Coulomb spectrum are however not extremely sensitive to the value of C once it is large enough. Using $C = -1400\text{MeV}$ instead would only reduce the nonperturbative contributions for the ground and first excited bottomonium-states in table 1 to 6 and 75MeV respectively.

The fact that a perturbative evaluation in ϕ is not appropriate in (33) can also not be circumvented by including loop corrections (higher orders in α_s than we have treated so far) to the perturbative Coulomb potential. Titard and Ynduráin[27] recently proposed to modify the perturbative part of the interaction in the following manner

$$\frac{\alpha_s}{r} \rightarrow \frac{\alpha_s(\mu^2) [1 + (a_1 + \gamma_E \beta_0/2) \alpha_s(\mu^2)/\pi]}{r} + \frac{\beta_0 \alpha_s^2(\mu^2) \log r \mu}{2\pi r}, \quad (34)$$

where the appropriate constants for the SU(3) color group with 4 light quark flavors are $\beta_0 = 8.33$ and $a_1 = 1.47$. The first term of (34) which contains a piece of one-loop radiative corrections was taken by Titard and Ynduráin as an effective Coulomb potential and solved exactly. The effective coupling constant is defined as

$$\tilde{\alpha}_s(\mu^2) = [1 + (a_1 + \gamma_E \beta_0/2) \frac{\alpha_s(\mu^2)}{\pi}] \alpha_s(\mu^2). \quad (35)$$

The second term in (34) was treated to first order in perturbation theory. A new scale parameter μ was introduced which depends on the quarkonium system under consideration. Taking the effective Coulomb potential we obtain the deviation of the eigenvalues from 2nd-order perturbation theory in ϕ shown in table 2 for two sets of parameters in the bottomonium system (still neglecting $1/m$ corrections).

The deviation of the exact correction from the perturbative estimate is reduced somewhat, especially in the second case, but still far from negligible. In assessing the quality of a perturbative evaluation in this case, one should also keep in mind that the scale parameter μ of Ref.[27] was chosen in such a way that the 2nd order correction in ϕ is precisely canceled by the correction terms to the effective Coulomb potential in (34). For the first set

of parameters this cancellation occurs for the ground state energy. The second set was chosen so that the splitting between the first excited state and the ground state is not affected to 2nd order. This is an arbitrary procedure which requires an additional parameter and does not cure the problem that perturbation theory simply does not apply (as table 2 clearly indicates).

We therefore will not include these modifications to the Coulomb force in our discussion of the numerical solution to the full set of coupled equations (30). The eigenvalues we obtained at order $(1/m)^0$ are summarized in table 3 and compared to those obtained by Eichten et al.[16] with the phenomenological funnel potential. Note that we only found 3 or 4 stable solutions in the charmonium and bottomonium systems respectively. We used the same values for the quark masses and the coupling constant as Eichten et al.[16]. The constant C was chosen to reproduce the experimental splitting (not confirmed[26]) between η_c and η'_c . It was not adjusted in the bottomonium system. The gluon condensate value $\phi^2 = (360\text{MeV})^4$ that we used is within QCD-sumrule estimates[28] for this nonperturbative quantity. All eigenvalues were finally shifted by $E_0 = 98\text{MeV}$ to give the correct η_c mass. (This small shift in the overall energy normalization can be eliminated by a slight change of $\sim 50\text{MeV}$ in the quark masses used by Eichten et al.[16] and a corresponding small adjustment of the other parameters. To have a more direct comparison of the wavefunctions and spectra, we refrained from making these adjustments here.)

The inclusion of $1/m$ potential terms lifts the degeneracy of pseudoscalar- and vector- quarkonia and we restrict our discussion here to the pseudoscalar case. The B and E_2 -states now couple in, but generally have small norms because the coupling is of order $1/m$. The effects of the $1/m$ corrections in the splittings are shown in table 4.

We also show in table 4 the results obtained using a regularized running coupling constant with the correct two-loop behaviour for small distances[29]:

$$\alpha_s = \alpha_s(r) = 4\pi \frac{1}{b_0 f(r)} \left[1 + \frac{2\gamma_E + 53/75}{f(r)} - \frac{462 \ln(f(r))}{625 f(r)} \right]$$

where

$$f(r) = \ln \left[\frac{1}{(\Lambda r)^2} + b \right]$$

$$b_0 = 25/3$$

In the above expression $\gamma_E = 0.5772$, and as in [29], we used $\Lambda = 300MeV$ and $b = 19$.

In Figs. 2 and 3 we show the eigenfunctions for the various components of our quarkonium states (7) as well as the corresponding eigenfunction of Eichten et al.[16]. The singlet component of our ground state wave functions are very similar to those of the funnel potential. At small radii this is true also for the excited states, since the coupling to “octet” components is proportional to r in (30). The “octet”-components increase with increasing excitation energy of the quarkonium and lead to the appearance of extra nodes in the higher lying “singlet” wave functions at large radii (since this is a coupled channel problem, the extra nodes do not mean that we missed some bound states). As noted earlier, the E -components of the meson state contribute negatively to its norm. All the stable quarkonia states that we found are however positive norm states. We could not obtain any stable state where the octet components are dominant.

Let us speculate at this point on the fact that only very few stable quarkonium states were found. This is of course quite in line with the experimental observation that only a few heavy quarkonia are stable against decay by strong interactions. For reasons which we had not anticipated, this basic property seems already to be incorporated in the nonhermitian coupling to E -components.

The strength of this coupling in our model is however determined by the gluon condensate ϕ^2 , which is not expected to vanish even in the purely gluonic theory. From table 3 we see that the instability sets in at an excitation energy of between 1 – 1.3GeV in this model. Since we cannot account for the decay into light $q\bar{q}$ -mesons with a parameter which is essentially independent of the number of (light) flavours, we speculate that the nonhermitian coupling proportional to ϕ effectively accounts for the possible decay channel

$$\text{Quarkonium}^* \longrightarrow \text{Quarkonium} + \text{Gluonium}, \quad (36)$$

in this model. From the fact that we do not seem to find any stable quarkonium state more than 1.3GeV above the ground state, we would estimate this to be the threshold for the production of the lightest gluonium. This estimate of the lightest gluonium mass $m_G \gtrsim 1.3GeV$, is in almost perfect agreement with our previous interpretation of the energy shift $-C = m_g \sim m_G/2$.

Since the production of light $q\bar{q}$ -pairs has a much lower threshold, it is this process which limits the stability of physical quarkonia. We therefore

expect this model, which does not (not even effectively) incorporate this decay channel, to still predict more stable states than are experimentally observed.

The influence of a running constant is more visible in the wave functions at the origin, which are important for production and annihilation decays. We compare the wave functions (not multiplied by r or r^2) with and without running coupling constants in figures 4 and 5.

6 Conclusion

We developed a hamiltonian formalism, which enabled us to estimate the effect of a nonperturbative gluonic ground state on quarkonia in a systematic short distance, weak coupling and $1/m$ expansion of the effective hamiltonian. The gauge invariant basis[8] was extended to include color octet quark-antiquark pairs coupled to vacuum fluctuations. Hamiltonian matrix elements in this basis are gauge invariant to the order considered in the short distance expansion.

After separating hard and soft gluons in the gauge (13), we obtained the effective hamiltonian neglecting radiative corrections to the Coulomb interaction from hard gluons. We thus neglected the logarithmic corrections to the effective coupling strength at very short distances. The correct behaviour of the potential for $r < 0.2fm$ can in principle be included by “hand” in a modification of the Coulomb part of the interaction[19]. Although the quarkonium spectrum is not very sensitive to this correction at small distances, it could become important for the evaluation of decay widths, which depend on the wave function at the origin.

In the limit of very heavy quark masses, where all terms of order $1/m$ can be neglected, an energy independent effective potential was obtained, which shows an approximately linear behaviour at intermediate distances $0.4fm < r < 0.8fm$ with an effective string tension of $\sim 840MeV$. In our approach this behaviour of the potential arises due to the nonperturbative structure of the gluonic vacuum parametrized by its gluon condensate and was not assumed from the outset as in most of phenomenological quark models. This potential compares favourably with recent lattice results[25], the discrepancies at very small $r < 0.2fm$ being due to our neglect of radiative corrections. Our short distance expansion for the effective potential however

is only valid for $r < 0.9fm$, beyond which the potential acquires an imaginary part. An extension of the model to larger distances would require a more detailed knowledge of the vacuum structure in the form of higher dimensional condensates, or some other effective parametrization of this structure. The approach in this case would become increasingly phenomenological and also more complicated in this case. Its predictive power is therefore probably limited to heavy quarkonia, where a detailed knowledge of the potential for very large radii does not seem to be necessary.

We showed that the numerical solution of the coupled channel problem for vector- and pseudoscalar- quarkonia (they are degenerate to order $1/m$) is feasible and an exact diagonalization of the hamiltonian in the extended basis therefore possible. The resulting exact spectrum does not show the far too rapid increase of the eigenvalues with the principal quantum number of the perturbative approach to the vacuum effects[9][20]. In addition to the usual “singlet” wave-functions describing the quark and anti-quark of the quarkonium when they are coupled in a colour singlet, we also obtain the “octet” components of quarkonium states describing the quarks in the octet configuration when an additional (soft) gluon is around. Since hadronic decays may be very sensitive to this “octet” configuration with the creation of an additional octet $q\bar{q}$ -pair from a hard gluon, this approach opens the possibility of estimating nonperturbative contributions to hadronic decays.

We compare our results for the spectrum and wavefunctions with those of the Cornell potential[16] for pseudoscalar quarkonia. With the standard value for the gluon condensate and quark masses and coupling constant used by the Cornell group we obtain the correct splitting between η_c and η'_c and make predictions for the η_b 's. Our main concern was however a better theoretical understanding and justification of the phenomenological ingredients common to most nonrelativistic models for heavy quarkonia and we refrained from adjusting the few parameters of this approach to optimally reproduce the experimental data. A better description of the potential at short distances with the inclusion of radiative corrections to the Coulomb force and the consideration of hadronic decay channels is clearly desirable before a detailed comparison with phenomenology is attempted.

Acknowledgements

We thank K. Langfeld for help in the numerical solution of the coupled equations and for useful discussions. One of the authors (C.A.A.N.) is grateful for the kind hospitality extended to him at the Institut für Theoretische Physik der Technischen Universität München. He would also like to thank Professors H. Leutwyler and K. Yazaki for helpful discussions. M.S. would like to thank St.Glazek for suggesting the extension of the basis. C.A.A.N. is supported by FAPESP/Brazil. F.S.N. is supported by CNPq and FAPESP/Brazil. M.S. is supported by Deutsche Forschungsgemeinschaft with grant Scha/1-2.

References

- [1] G.T. Bodwin, E. Braaten and G.P. Lepage, Phys.Rev. D 46 (1992) R1914; G.T. Bodwin, E. Braaten and G.P. Lepage, Phys.Rev. D 51 (1995) 1125.
- [2] W.E. Caswell and G.P. Lepage, Phys.Lett. B 167 (1986) 437.
- [3] P. Cho and A.K. Leibovich, Phys. Rev. D 53 (1995) 150; P. Cho and A.K. Leibovich, CALT-68-2026 ; M.L. Mangano and A. Petrelli, Phys. Lett. B 352 (1995) 445.
- [4] CDF Collaboration, “ J/Ψ , $\Psi' \rightarrow \mu^+\mu^-$ and $B \rightarrow J/\Psi, \Psi'$ Cross Sections,” FERMILAB-CONF-94/136-E, contribution GLS0208 to the 1994 International Conference on High Energy Physics, Glasgow.
- [5] E. Braaten and T.C. Yuan, Phys.Rev.Lett. 71 (1993) 1673; C.-H. Chang and Yu-Qi Chen, Phys.Lett b 284 (1991) 127; Phys.Rev. D 46 (1992) 3845.
- [6] E. Braaten, M.A. Doncheski, S. Fleming and M. Mangano, Phys.Lett. B 333 (1994) 548; M. Cacciari and M. Greco, Phys.Rev.Lett. 73 (1994) 1586.
- [7] E. Braaten and S. Fleming, Phys.Rev.Lett. 74 (1995) 3327.
- [8] S. Glazek and M. Schaden, Phys.Lett. B 198(1987) 42.

- [9] M.B. Voloshin, Nucl.Phys. B 154 (1979) 365.
- [10] M.B. Voloshin, Sov.J.Nucl.Phys. 36 (1982) 143.
- [11] L.L. Foldy and S.A. Wouthuysen, Phys.Rev. 78 (1950) 29; J.G. Körner and G. Thompson, Phys.Lett. B 264 (1991) 185; A. Das and V.S. Mathur, Phys.Rev. D 49 (1994) 2508; A. Das, preprint UR-1325 (1993).
- [12] G. 't Hooft, Nucl.Phys. B 62 (1973) 444; J. Honerkamp, Nucl.Phys. B 36 (1972) 130; B.S. DeWitt, Phys.Rev. 162 (1967) 1195; 1239.
- [13] G. Curci, A.Di Giacomo, G. Paffuti, Z.Phys. C 18 (1983) 135.
- [14] C.A.A. Nunes, PhD.Thesis, T.U.Munich (1992).
- [15] C. Callan, R. Dashen, and D. Gross, Phys.Rev. D 17 (1978) 2717.
- [16] E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.-M. Yan, Phys.Rev. D 17 (1978) 3090; Phys.Rev. D 21 (1980) 203.
- [17] F.S. Navarra and C.A.A. Nunes, Phys.Lett. B 356 (1995) 439.
- [18] M. Shifman, A. Vainshtein and V. Zakharov, Nucl.Phys. B 147 (1979) 385; 448.
- [19] W. Buchmüller and S.H.H. Tye, Phys.Rev. D 24 (1981) 132.
- [20] H. Leutwyler, Phys.Lett. 98 B (1981) 447.
- [21] C. Quigg and J.L. Rosner, Phys.Lett. B 71 (1977) 153; Phys.Rep. 56 (1979) 167; A. Martin, Phys.Lett. 93 B (1980) 338; Phys.Lett. B 100 (1981) 511.
- [22] J.C. Collins, A. Duncan, S.D. Joglekar, Phys.Rev. D 16 (1977) 438; S.L. Adler, J.C. Collins, A. Duncan, Phys.Rev. D 15 (1977) 1712; N.K. Nielsen, Nucl.Phys. B 120 (1980) 212.
- [23] H. Leutwyler, Nucl.Phys. B 179 (1981) 129.
- [24] H. Chen, J. Sexton, A. Vaccarino, D. Weingarten, Nucl.Phys. B(Suppl.)34 (1994) 357.

- [25] K.Schilling,G.S. Bali, Nucl.Phys.B (Suppl) 34 (1994) 147.
- [26] L.Montanet et al., Phys.Rev. D 50 (1994) 1.
- [27] S. Titard and F.J. Ynduráin, Phys.Rev. D 49 (1994) 6007; preprint FTUAM 94-6.
- [28] L.J. Reinders, H.R. Rubinstein and S. Yasaki, Phys.Rep. 127 (1986) 1.
- [29] K. Igi and S. Ono, Phys.Rev. D 33 (1986) 3349.

Figure 1: (a) Effective potential (solid curve) from (32) and Coulomb potential (dotted curve) with $\alpha_s = 0.39$, $C = -756$ MeV and $\phi^2 = (360 \text{ MeV})^4$. (b) Effective potential (solid curve) with the same parameters as in (a). The potential extracted from lattice data[25] with $\sqrt{\sigma} = 365$ MeV (dot-dashed curve) and $\sqrt{\sigma} = 505$ MeV (dashed curve). Cornell potential[16] (dotted curve) with $\alpha_s = 0.39$ and $a = 2.34 \text{ GeV}^{-1}$.

Figure 2: The wavefunctions of the ground and 1st excited pseudoscalar states of charmonium are shown in figures (a) and (b) respectively. The solid curve is the wavefunction for the funnel potential[16] for comparison. The long dashed curve is the singlet component $S(r)$ of the quarkonium state in our calculation. The dashed, dotted and dot-dashed curves are the “octet” components $E_1(r)$, $E_2(r)$ and $B(r)$ of equation (30) respectively. The solution was obtained with the parameters $m_c = 1840$ MeV, $\alpha_s = 0.39$, $\phi^2 = (360 \text{ MeV})^4$ and $C = -756$ MeV.

Figure 3: The wavefunctions of the ground, 1st- and 2nd excited pseudoscalar states of bottomonium are shown in figures (a), (b) and (c) respectively. The solid curve is the wavefunction for the funnel potential[16] for comparison. The long dashed curve is the singlet component $S(r)$ of the quarkonium state in our calculation. The dashed, dotted and dot-dashed curves are the “octet” components $E_1(r)$, $E_2(r)$ and $B(r)$ of equation (30) respectively. The solution was obtained with the parameters $m_b = 5170$ MeV, $\alpha_s = 0.39$, $\phi^2 = (360 \text{ MeV})^4$ and $C = -756$ MeV.

Figure 4: The wavefunctions (not multiplied by r or r^2) of the ground and 1st excited pseudoscalar states of charmonium are shown in figures (a) and (b) respectively. The solid curves are for the results without a running coupling constant. The dashed curves are obtained with a running coupling constant. The solutions were obtained with the parameters $m_c = 1840$ MeV, $\alpha_s = 0.39$ (without running), $\Lambda = 300$ MeV and $b = 19$ (with running), $\phi^2 = (360 \text{ MeV})^4$ and $C = -756$ MeV.

Figure 5: The wavefunctions of the ground, 1st- and 2nd excited pseudoscalar states of bottomonium are shown in figures (a), (b) and (c) respectively. The solid curves are for the results without a running coupling constant. The dashed curves are obtained with a running coupling constant. The solutions were obtained with the parameters $m_b = 5170$ MeV, $\alpha_s = 0.39$ (without running), $\Lambda = 300$ MeV and $b = 19$ (with running), $\phi^2 = (360 \text{ MeV})^4$ and $C = -756$ MeV.

Table 1: The Coulomb energy in MeV is presented in the first column for the ground state and first excitation of $c\bar{c}$ and $b\bar{b}$ (pseudoscalar or vector). Second and third columns contain the nonperturbative contributions in MeV calculated respectively within perturbation theory and with our method. We used $m_c = 1840$ MeV, $m_b = 5170$ MeV, $\alpha_s = 0.39$, $\phi^2 = 0.012 \text{ GeV}^4$ and $C = -756$ MeV.

	Coulomb	Pert. theory	Non-pert.
$\eta_c, J/\psi$	-124.4	311	66
η'_c, ψ'	-31.1	21525	424
η_b, Υ	-349.5	14	8
η'_b, Υ'	-87.4	970	111

Table 2: The Coulomb energy in MeV is presented in the first column for the ground state and first excitation of $b\bar{b}$ (pseudoscalar or vector) at two different scales μ . Second and third columns contain the nonperturbative contributions in MeV calculated respectively within perturbation theory and with our method. We used $C = -756$ MeV, $\phi^2 = 0.042 \text{ GeV}^4$ and for $\mu = 1.44 \text{ GeV}$: $m_b = 4866$ MeV, $\bar{\alpha}_s = 0.38$. For $\mu = 0.99 \text{ GeV}$: $m_b = 5010$ MeV, $\bar{\alpha}_s = 0.54$.

	Coulomb	Pert. theory	Non-pert.
η_b, Υ ($\mu = 1.44 \text{ GeV}$)	-312	25	11
η'_b, Υ' ($\mu = 1.44 \text{ GeV}$)	-78	1762	129
η_b, Υ ($\mu = 0.99 \text{ GeV}$)	-649	6	3
η'_b, Υ' ($\mu = 0.99 \text{ GeV}$)	-162	396	72

Table 3: Masses in MeV of pseudoscalar quarkonia with the funnel potential and with our effective nonperturbative hamiltonian. Parameters used: $m_c = 1840$ MeV, $m_b = 5170$ MeV, $\alpha_s = 0.39$, $\phi^2 = (360 \text{ MeV})^4$, $C = -756$ MeV.

	funnel	nonpert.
η_c	2980	2980
η'_c	3571	3594
η''_c	3994	3993
η_b	9213	9344
η'_b	9805	9739
η''_b	10150	10084
η'''_b	10427	10610

Table 4: Energy splittings among radial excitations of pseudoscalar charmonium and bottomonium. The parameters are the same as in table 3. For the running coupling constant (see text) we use $\Lambda = 300$ MeV and $b = 19$. First column: no $1/m$ terms. Second column: $1/m$ corrections included. Third column: $1/m$ terms and running coupling constant.

	no $1/m$	$1/m$ included	$1/m$ and running
$c\bar{c}$ - 2S-1S	614	666	593
$b\bar{b}$ - 2S-1S	395	397	413
$b\bar{b}$ - 3S-1S	740	753	738





