



Instituto de Física
Universidade de São Paulo

Hyperon-nucleon coupling from QCD sum rules

Bracco, Me^a; Navarra, FSb; Nielsen, M^b

*^aInstituto de Física, Universidade do Estado do Rio de Janeiro,
Rua São Francisco Xavier 524 20559-900, Rio de Janeiro, RJ, Brasil*

*^bInstituto de Física Universidade de São Paulo
C.P. 66318, CEP 05315-970, São Paulo, SP, Brasil*

Publicação IF - 1421/2000

1421

High Energy Physics - Phenomenology, abstract hep-ph/9908452

SBI/IFUSP

BASE: 04

SYS N°: 1097289

From: Marina Nielsen <mnielsen@if.usp.br>
Date: Tue, 24 Aug 1999 13:29:22 GMT (23kb)

Hyperon-nucleon coupling from QCD sum rules

Authors: M. E. Bracco, F. S. Navarra, M. Nielsen

Comments: 4 pages, 2 figures, espcrc2.sty included. Talk presented at QCD99, Montpellier, France (to appear in Nucl.Phys.B Proc. Suppl.)

The NKY coupling constant for $Y = \Lambda$ and Σ is evaluated in a QCD sum rule calculation. We discuss and extend the result of a previous analysis in the γ_5 structure and compare it with the result obtained with the use of the $\gamma_5 \sigma_{\mu\nu}$ structure. We find a huge violation of the SU(3) symmetry in the $\gamma_5 \sigma_{\mu\nu}$ structure.

Paper: Source (23kb), PostScript, or Other formats

(N.B.: delivery types and potential problems)

refers to , cited by

Links to: xxx, hep-ph, /find, /abs (-/+), /9908, ? form interface

www-admin@xxx.lanl.gov

Hyperon-nucleon coupling from QCD sum rules

M.E. Bracco ^a, F.S. Navarra ^b and M. Nielsen ^b

^aInstituto de Física, Universidade do Estado do Rio de Janeiro,
Rua São Francisco Xavier 524 - 20559-900, Rio de Janeiro, RJ, Brazil

^bInstituto de Física, Universidade de São Paulo,
C.P. 66318, 05389-970 São Paulo, SP, Brazil

The NKY coupling constant for $Y = \Lambda$ and Σ is evaluated in a QCD sum rule calculation. We discuss and extend the result of a previous analysis in the $\not{q}\gamma_5$ structure and compare it with the result obtained with the use of the $\gamma_5\sigma_{\mu\nu}$ structure. We find a huge violation of the $SU(3)$ symmetry in the $\gamma_5\sigma_{\mu\nu}$ structure.

In understanding the dynamics of kaon-nucleon scattering or the strangeness content of the nucleon using hadronic models, it is important to know the hadronic coupling constants involving the kaons. Among them, $g_{NK\Lambda}$ and $g_{NK\Sigma}$ are the most relevant coupling constants. To determine these couplings using the QCD sum rules [1] one can follow two different approaches: a) the two-point function, where the nucleon and hyperon fields are sandwiched between the vacuum and kaon states, or b) the three-point function where three interpolating fields are sandwiched between vacuum states.

In the case of the pion-nucleon coupling constant, in a pioneer calculation [2] both approaches showed to reproduce the phenomenological value fairly well. However, in this first study the continuum contribution was neglected and since then many calculations were done including higher order terms in the operator product expansion (OPE) and the continuum contribution [3], going beyond the soft-pion limit and including also pole-continuum transitions [4].

For the nucleon-kaon-hyperon coupling constant there are also QCD sum rules calculations based on the two- and three-point functions [5-7]. The advantages of the three-point function calculation is that it allows for the calculation of the form factors at the hadronic vertices.

In the strange sector, the nucleon-kaon-hyperon form factors are used, for instance, to evaluate the strange radius of the nucleon using

the kaon cloud [8] and, therefore, a theoretically founded evaluation of these form factors is welcome.

We will calculate the g_{NKY} coupling constant using the three-point function

$$A(p, p', q) = \int d^4x d^4y e^{ip'x} e^{-iqy} \times \langle 0 | T \{ \eta_Y(x) j_5(y) \bar{\eta}_N(0) \} | 0 \rangle \quad (1)$$

where $j_5 = \bar{s}i\gamma_5 u$.

As it is well known from two-point sum rules for baryons, there is a continuum of choices for the baryon interpolating fields. Of course the results should be independent of the choice of the current, if we considered an infinity number of terms in the OPE and if we had a perfect model for the continuum contribution in the phenomenological side. However, the OPE has to be truncated and we work with a very simple model for the continuum contribution. Therefore, the results do depend on the choice of the currents. For the proton Λ and Σ we can write general currents as [9]

$$\eta_P = 2\varepsilon_{abc} [(u_a^T C d_b) \gamma_5 u_c + b(u_a^T C \gamma_5 d_b) u_c], \quad (2)$$

$$\eta_Y = 2[\eta_{Y_1} + b\eta_{Y_2}], \quad (3)$$

where b is a parameter and

$$\eta_{\Lambda_1} = \frac{1}{\sqrt{6}} \varepsilon_{abc} [2(u_a^T C d_b) \gamma_5 s_c + (u_a^T C s_b) \gamma_5 d_c - (d_a^T C s_b) \gamma_5 u_c], \quad (4)$$