



**Instituto de Física
Universidade de São Paulo**

On the action principle for a system of
differential equations

D.M. Gitman, V.G. Kupriyanov

Publicação IF – 1638/2007

UNIVERSIDADE DE SÃO PAULO
Instituto de Física
Cidade Universitária
Caixa Postal 66.318
05315-970 - São Paulo - Brasil

On the action principle for a system of differential equations

D.M. Gitman¹, V.G. Kupriyanov^{1,2}

¹Instituto de Física, Universidade de São Paulo,
Caixa Postal 66318-CEP, 05315-970 São Paulo, S.P., Brazil

² Physics Department, Tomsk State University, Tomsk, 634050, Russia
E-mail: gitman@dfn.if.usp.br (D.M.Gitman), kvg@dfn.if.usp.br (V.G.Kupriyanov).

July 2, 2007

Abstract

We consider the problem of constructing an action functional for physical systems whose classical equations of motion cannot be directly identified with Euler-Lagrange equations for an action principle. Two ways of action principle construction are presented. From simple consideration, we derive necessary and sufficient conditions for the existence of a multiplier matrix which can endow a prescribed set of second-order differential equations with the structure of Euler-Lagrange equations. An explicit form of the action is constructed in case if such a multiplier exists. If a given set of differential equations cannot be derived from an action principle, one can reformulate such a set in an equivalent first-order form which can always be treated as the Euler-Lagrange equations of a certain action. We construct such an action explicitly. There exists an ambiguity (not reduced to a total time derivative) in associating a Lagrange function with a given set of equations. We present a complete description of this ambiguity. The general procedure is illustrated by several examples.

1 Introduction

The problem of constructing an action functional for a given set of differential equations is known in literature as the inverse problem of the calculus of variations for the Newtonian mechanics. In its classical setting [1] consists in solving of the variational equation

$$\frac{\delta S[q]}{\delta q^i(t)} = g_i, \quad (1)$$

where $g_i(t, q^i, \dot{q}^i, \dots) = 0$ is some given system of differential equations with respect to unknown functions $q^i(t)$, and $S[q]$ is a local functional to be determined. The condition of locality requires the existence of a function $L(t, q, \dot{q}, \dots)$ (Lagrangian), such that the functional $S[q]$ (action) would be written as an integral

$$S[q] = \int dt L. \quad (2)$$

In other words, the essence of the inverse problem of the calculus of variations consists in finding a variation principle for a given system of differential equations. This problem has been under consideration for more than a hundred years. As early as 1887 Helmholtz [1] presented a criterion of commutativity for second variational derivatives from which immediately follows the necessary (and with some restrictions also sufficient) condition of solvability of the equation (1):

$$\frac{\delta g_i(t)}{\delta q^j(s)} = \frac{\delta g_j(s)}{\delta q^i(t)}. \quad (3)$$

If this condition holds, the system $g_i(t, q^i, \dot{q}^i, \dots) = 0$ is called Lagrangian system, if not the system is non-Lagrangian. In 1894 Darboux [2] solved the problem for one dimensional case. In 1941 the case of two degrees of freedom was investigated by Douglas [3]; in particular, he presented examples of second-order equations which

cannot be obtained from the variational principle. Afterwards many authors (see e.g., [6]-[13] and references therein) investigated this problem for multidimensional systems.

In the present work we consider the question of the construction an action principle for a given system of differential equations using integrating multiplier method [3]-[9]. The integrating multiplier is such a nonsingular matrix which being multiplied by a given set of differential equations reduces this set to a standard Euler-Lagrange form. In section 2 we present a simple derivation for necessary and sufficient conditions for an integrating multiplier for a system of second-order equations. We also construct the explicit form of Lagrangian in a case an integrating multiplier does exist. Then we apply our method for investigating the inverse problem to some simple models. In particular, we construct an action principle for multidimensional dissipative systems. We also consider an example of a linear dynamical system whose equations of motion do not admit an integrating multiplier, and, as a consequence, cannot be obtained from the minimum action principle.

Note that it is always possible to reduce non-Lagrangian second-order equations of motion to an equivalent set of first-order differential equations. From Helmholtz criterion (3) we find necessary and sufficient conditions for an integrating multiplier for such equations. It turns out that in the first-order formalism an integrating multiplier is always exist and can be constructed by the solution of the Cauchy problem for the equations in question, the corresponding consideration is represented in section 3 and, partially, is based on results of works [10] and [13]. Then we construct the action functional explicitly. Thus, we show that systems traditionally called non-Lagrangian ones are, in fact, equivalent to some first-order Lagrangian systems. As an example, we construct a first-order action functional for any linear dynamical system.

2 Action functional for a set of second-order equations

2.1 General consideration

Let a system with n degrees of freedom be described by a set of n second-order differential equations of motion, solvable with respect to second-order time derivatives. Suppose such a set has a form

$$\ddot{q}^i - f^i(t, q, \dot{q}) = 0, \quad i = 1 \dots n, \quad (4)$$

where $f^i(t, q, \dot{q})$ are some functions of the indicated arguments, and by dots above we denote time derivatives of the coordinates. Let us construct an action principle for this set. If (4) cannot be directly identified with Euler-Lagrange equations, one can find an integrating multiplier, i.e., such a nonsingular matrix $h_{ij}(t, q, \dot{q})$ that being multiplied by (4)

$$h_{ij} [\ddot{q}^j - f^j(t, q, \dot{q})] = 0 \quad (5)$$

reduces this set to a standard Euler-Lagrange form for some Lagrangian $L(t, q, \dot{q})$,

$$\frac{\partial L}{\partial q^i} - \frac{\partial^2 L}{\partial t \partial \dot{q}^i} - \frac{\partial^2 L}{\partial \dot{q}^i \partial \dot{q}^j} \dot{q}^j - \frac{\partial^2 L}{\partial \dot{q}^i \partial \dot{q}^j} \ddot{q}^j = 0. \quad (6)$$

In order to identify (5) with (6) we need to ensure that

$$\frac{\partial^2 L}{\partial \dot{q}^i \partial \dot{q}^j} = h_{ij}, \quad (7)$$

$$\frac{\partial L}{\partial q^i} - \frac{\partial^2 L}{\partial t \partial \dot{q}^i} - \frac{\partial^2 L}{\partial \dot{q}^i \partial \dot{q}^j} \dot{q}^j = h_{ij} f^j. \quad (8)$$

Provided that an integrating multiplier is known, equations (7)-(8) can be interpreted as a system of equations for a Lagrange function L . We are going to solve the set of equations (7)-(8). Its consistency conditions will give us all the necessary and sufficient conditions for an integrating multiplier. Assuming that L is a smooth function of the indicated arguments, the consistency condition for equation (7) imply that

$$h_{ij} = h_{ji}, \quad \frac{\partial h_{ij}}{\partial \dot{q}^k} = \frac{\partial h_{kj}}{\partial \dot{q}^i}. \quad (9)$$

If (9) does hold one can solve equation (7). To this end, we remind that the general solution of the equation $\partial f / \partial q^i = g_i$, provided the vector g_i is a gradient, is

$$f(q) = \int_0^1 ds q^i g_i(sq) + c,$$

where c is a constant. Taking the above fact into account, we obtain for L (we do not consider global problems which can arise from non-trivial topology of the configuration space) the following representation:

$$L = K(t, q, \dot{q}) + l_i(t, q)\dot{q}^i + l_0(t, q), \quad (10)$$

where

$$K(t, q, \dot{q}) = \int_0^1 da \dot{q}^j \left[\int_0^1 db \dot{q}_1^i h_{ij}(t, q, b\dot{q}_1) \right]_{\dot{q}_1 = a\dot{q}} \quad (11)$$

and $l_0(t, q)$, $l_i(t, q)$ are some functions of the indicated arguments. To find these functions, we use equation (8). Substituting (10) into (8), we get

$$\frac{\partial K}{\partial q^i} - \frac{\partial^2 K}{\partial \dot{q}^i \partial t} - \frac{\partial^2 K}{\partial \dot{q}^i \partial \dot{q}^j} \dot{q}^j + \left(\frac{\partial l_j}{\partial q^i} - \frac{\partial l_i}{\partial q^j} \right) \dot{q}^j - \frac{\partial l_i}{\partial t} + \frac{\partial l_0}{\partial q^i} = h_{ij} f^j. \quad (12)$$

Differentiating this equation over \dot{q}^k , we obtain:

$$\frac{\partial l_k}{\partial q^i} - \frac{\partial l_i}{\partial q^k} = L_{ik}, \quad (13)$$

where

$$L_{ik} = \frac{\partial^2 K}{\partial \dot{q}^i \partial \dot{q}^j} - \frac{\partial^2 K}{\partial \dot{q}^j \partial \dot{q}^i} + \frac{\partial h_{ik}}{\partial t} + \dot{q}^j \frac{\partial h_{ik}}{\partial \dot{q}^j} + \frac{\partial}{\partial \dot{q}^k} (h_{ij} f^j). \quad (14)$$

This equation is a differential equation for l_i . The consistency conditions of equation (13) imply, that first of all, the symmetric part of L_{ik} is zero, which can be written as

$$\hat{D}h_{ik} + \frac{1}{2} \left(h_{ij} \frac{\partial f^j}{\partial \dot{q}^k} + h_{kj} \frac{\partial f^j}{\partial \dot{q}^i} \right) = 0, \quad (15)$$

where

$$\hat{D} = \frac{\partial}{\partial t} + \dot{q}^j \frac{\partial}{\partial q^j} + f^j \frac{\partial}{\partial \dot{q}^j}.$$

Using (15), one can rewrite (14) as follows

$$L_{ik} = \frac{\partial^2 K}{\partial \dot{q}^i \partial \dot{q}^k} - \frac{\partial^2 K}{\partial \dot{q}^k \partial \dot{q}^i} + A_{ik}, \quad A_{ik} = \frac{1}{2} \left(h_{ij} \frac{\partial f^j}{\partial \dot{q}^k} - h_{kj} \frac{\partial f^j}{\partial \dot{q}^i} \right) \quad (16)$$

Next, L_{ik} does not depend on velocities, i.e., $\partial L_{ik} / \partial \dot{q}^l = 0$, which yields

$$\frac{\partial h_{kl}}{\partial q^i} - \frac{\partial h_{il}}{\partial q^k} = \frac{\partial}{\partial \dot{q}^l} A_{ik}. \quad (17)$$

And, finally, the Jacobi identity

$$\frac{\partial L_{ik}}{\partial q^l} + \frac{\partial L_{kl}}{\partial q^i} + \frac{\partial L_{li}}{\partial q^k} = 0 \Rightarrow \frac{\partial A_{ik}}{\partial q^l} + \frac{\partial A_{kl}}{\partial q^i} + \frac{\partial A_{li}}{\partial q^k} = 0. \quad (18)$$

Provided h_{ij} obeys equations (15), (17) and (18), l_i can be found from equation (13). We remind that the general solution for l_i of equation (13) is given by

$$l_i(t, q) = \int_0^1 da q^k L_{ki}(t, aq) + \frac{\partial \varphi(t, q)}{\partial q^i}, \quad (19)$$

where $\varphi(t, q)$ is an arbitrary function.

Now from equation (12) we can find l_0 ; to this end let us rewrite it as follows:

$$\frac{\partial l_0}{\partial q^i} = m_i, \quad (20)$$

where

$$m_i = h_{ij} f^j - \frac{\partial K}{\partial q^i} + \frac{\partial^2 K}{\partial t \partial \dot{q}^i} + \dot{q}^j \frac{\partial^2 K}{\partial q^j \partial \dot{q}^i} - \dot{q}^j L_{ij} + \frac{\partial l_i}{\partial t}. \quad (21)$$

The consistency conditions of (20) imply, that, first, m_i does not depend on velocities, i.e., $\partial m_i / \partial \dot{q}^k = 0$. This condition is provided by equation (15). And second, the vector m_i must be a gradient:

$$\frac{\partial m_i}{\partial q^k} - \frac{\partial m_k}{\partial q^i} = \frac{\partial A_{ik}}{\partial t} + \dot{q}^j \frac{\partial A_{ik}}{\partial q^j} + \frac{\partial}{\partial q^k} (h_{ij} f^j) - \frac{\partial}{\partial q^i} (h_{kj} f^j) = 0. \quad (22)$$

Taking into account (9), (15) and (17), one gets from (22) the following algebraic condition:

$$h_{ij} B_k^j - h_{kj} B_i^j = 0, \quad (23)$$

where

$$B_j^i = \frac{1}{2} \frac{\partial f^i}{\partial \dot{q}^m} \frac{\partial f^m}{\partial \dot{q}^j} - \dot{D} \frac{\partial f^i}{\partial \dot{q}^j} + 2 \frac{\partial f^i}{\partial q^j}.$$

If h_{ij} obeys (23), then from (20) one gets

$$l_0(t, q) = \int_0^1 da q^k m_k(t, aq) + \frac{\partial \varphi(t, q)}{\partial t} + c(t), \quad (24)$$

where $c(t)$ is an arbitrary function of time.

Thus, we have proved the following statement: *iff for a given set of second-order ordinary differential equations (4) there exist such a non-singular matrix $h_{ij}(t, q, \dot{q})$ that obeys equations (9), (15), (17), (18) and (23), then this set can be obtained from the variational principle with the Lagrangian (10), where functions $K(t, q, \dot{q})$, $l_i(t, q)$ and $l_0(t, q)$ are defined by (11), (19) and (24) correspondingly and the functions $\varphi(t, q)$ and $c(t)$ are arbitrary functions of the indicated arguments.*

The arbitrariness related to the functions $\varphi(t, q)$ and $c(t)$ enter Lagrangian (10) via the total time derivative of a function F ,

$$F = \varphi(t, q) + \int c(t) dt.$$

Note that an integrating multiplier h_{ij} , and as a consequence the Lagrange function L does exist, but however not for any set of equations (4). In Section 3 we consider an example of a dynamical system which does not admit the existence of an integrating multiplier. However, if it does exist, it is not unique [13]-[19], e.g., if the matrix h_{ij} is an integrating multiplier for a certain set (4), it is easy to see that the matrix $\tilde{h}_{ij} = c h_{ij}$, where $c \neq 0$ is a constant, is an integrating multiplier as well. Therefore, Lagrangian (10) leading to the set of equations (4) is not unique; for this set there exist as many inequivalent Lagrangians as integrating multipliers. Lagrangians corresponding to different integrating multipliers are known as s -equivalent Lagrangians.

In the one dimensional case $\ddot{q} - f(t, q, \dot{q}) = 0$, an integrating multiplier is a non-vanishing function $h(t, q, \dot{q})$ that obeys the equation

$$\frac{\partial h}{\partial t} + \dot{q} \frac{\partial h}{\partial q} + \frac{\partial}{\partial \dot{q}} (f h) = 0. \quad (25)$$

This is a first-order partial differential equation which obviously has a solution for any f and initial condition $h(t=0, q, \dot{q}) = h_0(q, \dot{q})$. As we can see, an answer to the question whether there exists a solution of the inverse problem of the calculus of variations depends on the number of degrees of freedom n . For $n = 1$ the answer is always positive, and there exist as many inequivalent Lagrangians as functions $h_0(q, \dot{q})$ of two variables. For $n \geq 2$ the answer is generally negative.

2.2 Examples

In this section we consider a possibility of constructing an action principle for some examples of dynamical systems. First of all, let us consider dissipative systems. Suppose we have an ideal system with the Lagrangian

$$L_0 = \frac{\dot{q}^2}{2} + V(q), \quad q = \{q^i\}, \quad i = 1 \dots n. \quad (26)$$

Let us consider the case when besides the potential conservative force $F^i = \frac{\partial V}{\partial q^i}$ there exist a friction force

$$F_{fric}^i = \alpha \dot{q}^i, \quad (27)$$

where α is a phenomenological friction coefficient which in general can depend on time. The equations of motion for such a system have the form

$$\ddot{q}^i = \frac{\partial V}{\partial q^i} + \alpha \dot{q}^i. \quad (28)$$

These equations are non-Lagrangian, but for this set it is possible to find an integrating multiplier. In the simplest case, when it does not depend on coordinates and velocities, it has the form

$$h_{ij} = e^{-2 \int \alpha dt} h_{ij}^0, \quad (29)$$

where h_{ij}^0 is an arbitrary, symmetric, nonsingular, constant matrix commuting with the matrix $V_{ij} = \partial^2 V / \partial q^i \partial q^j$. Using the statement of the previous section, we obtain the following Lagrangian:

$$L = \frac{1}{2} \dot{q}^i h_{ij} \dot{q}^j + \int_0^1 q^i h_{ij} \frac{\partial V(s\vec{q})}{\partial q^j} ds. \quad (30)$$

If one sets $h_{ij}^0 = \delta_{ij}$, Lagrangian (30) can be rewritten as

$$L = e^{-2 \int \alpha dt} L_0. \quad (31)$$

Note that once the friction coefficient goes to zero, Lagrangian (31) transforms into the initial Lagrangian (26).

Let us now consider the case when the potential in the initial Lagrangian is linear in velocities. For simplicity we consider the two-dimensional case

$$L_0 = \frac{1}{2} (\dot{x}^2 + \dot{y}^2 + \beta(\dot{x}y - \dot{y}x)), \quad (32)$$

Let us consider this system in the presence of dissipative the force (30). The equations of motion will have the form:

$$\begin{aligned} \ddot{x} &= \alpha \dot{x} - \beta \dot{y}, \\ \ddot{y} &= \beta \dot{x} + \alpha \dot{y}. \end{aligned} \quad (33)$$

As was shown in [12], this system describes a moving charged particle in a uniform magnetic field with radiation friction. In this case,

$$B_j^i = \begin{pmatrix} \alpha^2 - \beta^2 & -2\alpha\beta \\ 2\alpha\beta & \alpha^2 - \beta^2 \end{pmatrix}$$

and from equation (23) one immediately gets

$$tr(h_{ij}) = h_{11} + h_{22} = 0. \quad (34)$$

It is then easy to find that the general solution of the equations (9), (15), (17), (18) is defined by an arbitrary function $\phi(\zeta, \eta)$ and has a form

$$h_{ij} = \begin{pmatrix} F + \bar{F} & i(F - \bar{F}) \\ i(F - \bar{F}) & -(F + \bar{F}) \end{pmatrix}, \quad (35)$$

where $F = \phi(\dot{\xi} e^{-\gamma t}, \dot{\xi} - \alpha \xi) e^{-\gamma t}$, $\xi = x + iy$, $\gamma = \alpha + i\beta$ and the bar denotes a complex conjugation.

The simplest real solution can be found if we put $\phi = 1/\zeta$. We have

$$h_{ij} = \frac{2}{\dot{x}^2 + \dot{y}^2} \begin{pmatrix} \dot{x} & \dot{y} \\ \dot{y} & -\dot{x} \end{pmatrix}. \quad (36)$$

Using the formulas (10), (11), (19) and (24) we find the following Lagrangian:

$$L = \frac{1}{2} \dot{x} \ln(\dot{x}^2 + \dot{y}^2) + \dot{y} \arctan\left(\frac{\dot{x}}{\dot{y}}\right) + \alpha x - \beta y. \quad (37)$$

The corresponding Euler-Lagrange equations

$$\frac{\ddot{x}\dot{x} + \dot{y}\dot{y}}{\dot{x}^2 + \dot{y}^2} = \alpha, \quad \frac{\ddot{x}\dot{x} - \dot{y}\dot{y}}{\dot{x}^2 + \dot{y}^2} = \beta \quad (38)$$

are equivalent to the initial ones (33), with the exception of the point $\dot{x} = \dot{y} = 0$. Thus, we can see that in this case the inverse problem of the calculus of variations is solvable. Unfortunately, neither Lagrangian (37) nor any other Lagrangian constructed by the matrix (35) in the limit $\alpha \rightarrow 0$ transforms into the initial Lagrangian (32), modulo a total time derivative. This is because, according to the algebraic condition (34), the trace of the Hessian matrix of any Lagrangian for the set of equations (33) must be equal to zero and this property holds true after the limit $\alpha \rightarrow 0$ is taken. On the other hand, the trace of the Hessian matrix of the Lagrangian L_0 in (32) is equal to 2. This contradiction proves the statement.

Finally we consider example of dynamical system for which an integrating multiplier and, consequently, the possibility of the Lagrangian description does not exist. Douglas [3] showed that the set of second-order equations

$$\begin{aligned} \ddot{x} + \dot{y} &= 0, \\ \ddot{y} + x &= 0 \end{aligned}$$

does not admit an integrating multiplier. Let us make sure of this. To this end, let us assume the opposite, namely, let there exist such a non-degenerate matrix h_{ij} that obeys equations (9), (15), (17), (18) and (23). Then from the algebraic equation (23) it follows that h_{ij} must be diagonal ($h_{12} = h_{21} = 0$), since in this case

$$B_j^i = \begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix}.$$

Then, from condition (15) we immediately obtains that $h_{11} = 0$, and arrive at a contradiction, $\det h_{ij} = 0$.

Thus, we can see that an action functional in second-order formalism cannot be constructed not for some sets of differential equations. Nevertheless, as we show in the following section, it is always possible to construct an action principle for the equivalent set of first-order equations.

3 Action principle in the first-order form

Let a system with n degrees of freedom be described by a set of n non-Lagrangian second-order differential equations of motion. To construct an action principle, we replace these equations (which is always possible to do by introducing n additional variables, e.g. $p_i = \dot{q}^i$) by an equivalent set of $2n$ first-order differential equations, solvable with respect to time derivatives. Suppose such a set has a form

$$\begin{aligned} \dot{x}^\alpha &= f^\alpha(t, x), \\ x^\alpha &= (q^i, p_i), \quad \alpha = 1, \dots, 2n, \end{aligned} \quad (39)$$

where $f^\alpha(t, x)$ are some functions of the indicated arguments and by dots above we denote time derivatives of coordinates. Let us construct an action principle for this set. If (39) cannot be directly identified with Euler-Lagrange equations, then one can find an integrating multiplier, i.e. such a nonsingular matrix Ω^1 multiplication by which reduces initial set of differential equations (39) to a variational derivative:

$$g_\alpha [t] = \Omega_{\alpha\beta} (\dot{x}^\beta - f^\beta(t, x)) = \frac{\delta S}{\delta x^\alpha} = 0. \quad (40)$$

Since $g_\alpha [t]$ is a variational derivative it must obey the Helmholtz criterion [1]

$$\frac{\delta g_\alpha [t]}{\delta x^\beta (s)} = \frac{\delta g_\beta [s]}{\delta x^\alpha (t)}. \quad (41)$$

We will use this condition to find an integrating multiplier Ω . In the general case one can assume that Ω depends on time t , coordinates x^α and time derivatives up to the m order (m is a natural number), i.e. $g_\alpha [t] = g_\alpha (t, x, \dots, x^{(m)})$. Having in mind this form of g_α one rewrites (41) as

$$\sum_{i=0}^m \frac{\partial g_\alpha [t]}{\partial x^{\beta(i)}} \delta^{(i)} (t-s) = \sum_{j=0}^m \frac{\partial g_\beta [s]}{\partial x^{\alpha(j)}} \delta^{(j)} (s-t), \quad (42)$$

¹The integrating multiplier for the first-order equations we denote by Ω .

since

$$\frac{\delta}{\delta x^\beta(s)} x^{\alpha(k)}(t) = \left(\frac{d}{dt}\right)^k \frac{\delta x^\alpha(t)}{\delta x^\beta(s)} = \delta_\beta^\alpha \delta^{(k)}(t-s), \quad k = 0, 1, \dots$$

Differentiating the identity $f(t) \delta(t-s) = f(s) \delta(s-t)$ over t one finds

$$f(s) \delta^{(k)}(s-t) = (-1)^k \sum_{l=0}^k C_l^k f^{(l)}(t) \delta^{(k-l)}(t-s), \quad C_l^k = \frac{k!}{(k-l)!}. \quad (43)$$

Using (43) we rewrite (41) as

$$\sum_{i=0}^m \frac{\partial g_\alpha[t]}{\partial x^{\beta(i)}} \delta^{(i)}(t-s) = \sum_{j=0}^m (-1)^j \sum_{l=0}^j C_l^j \left[\left(\frac{d}{dt}\right)^l \frac{\partial g_\beta[t]}{\partial x^{\alpha(j)}} \right] \delta^{(j-l)}(t-s). \quad (44)$$

Comparing in (44) coefficient under $\delta^{(k)}(t-s)$ one gets equations on $g_\alpha(t, x, \dots, x^{(m)})$. When $k=0$ we have

$$\frac{\partial g_\alpha}{\partial x^\beta} - \sum_{j=0}^m (-1)^j \left(\frac{d}{dt}\right)^j \frac{\partial g_\beta}{\partial x^{\alpha(j)}} = 0. \quad (45)$$

Since $g_\alpha[t]$ depends only on derivatives up to the m order, coefficient under higher derivatives $x^{\alpha(2m)}$ in equation (45) must vanish. This coefficient is $(-1)^m \partial^2 g_\beta / \partial x^{\alpha(m)} \partial x^{\gamma(m)}$, which means that $g_\alpha[t]$ must be linear on the derivatives of the m order, i.e.

$$g_\alpha[t] = a_{\alpha\beta} \left(t, x, \dots, x^{(m-1)}\right) x^{\beta(m)} + b_\alpha \left(t, x, \dots, x^{(m-1)}\right),$$

where $a_{\alpha\beta}$ and b_α are some functions. Since Ω is nonsingular matrix, (40) should be a system of first-order equations, i.e. we have $m=1$ and $\Omega = \Omega(t, x)$.

Now comparing in (44) coefficient under $\delta^{(1)}(t-s)$ we obtain:

$$\Omega_{\alpha\beta} = -\Omega_{\beta\alpha}. \quad (46)$$

Then from (45) we have

$$\partial_\beta (\Omega_{\alpha\gamma} f^\gamma) - \partial_\alpha (\Omega_{\beta\gamma} f^\gamma) + \partial_t \Omega_{\alpha\beta} + \dot{x}^\gamma (\partial_\beta \Omega_{\alpha\gamma} - \partial_\alpha \Omega_{\beta\gamma} + \partial_\gamma \Omega_{\beta\alpha}) = 0.$$

Since Ω does not depend on \dot{x} one gets the following equations on Ω :

$$\partial_\alpha \Omega_{\beta\gamma} + \partial_\beta \Omega_{\gamma\alpha} + \partial_\gamma \Omega_{\alpha\beta} = 0 \quad (47a)$$

and

$$\partial_t \Omega_{\alpha\beta} + \mathcal{L}_f \Omega_{\alpha\beta} = 0, \quad (48)$$

where $\mathcal{L}_f \Omega_{\alpha\beta}$ is the Lie derivative of $\Omega_{\alpha\beta}$ along the vector field f^γ and $\partial_\alpha = \partial / \partial x^\alpha$, $\partial_t = \partial / \partial t$. Thus we see that for the set of first order equations an integrating multiplier is a nonsingular matrix which depends only on time t and coordinates x^α and obey the conditions (46), (47a) and (48).

Let us analyze equations (46)-(48) for the matrix $\Omega_{\alpha\beta}$ following our work [13]. It is known that the general solution $\Omega_{\alpha\beta}$ of equation (48) can be constructed with the help of a solution of the Cauchy problem for equations (39). Suppose that such a solution is known,

$$x^\alpha = \varphi^\alpha(t, x_{(0)}), \quad x_{(0)}^\alpha = \varphi^\alpha(0, x_{(0)}) \quad (49)$$

be a solution of equations (39) for any $x_{(0)} = \left(x_{(0)}^\alpha\right)$, and $\chi^\alpha(t, x)$ be the inverse function with respect to $\varphi^\alpha(t, x_{(0)})$, i.e.,

$$x^\alpha = \varphi^\alpha(t, x_{(0)}) \implies x_{(0)}^\alpha = \chi^\alpha(t, x), \quad x^\alpha \equiv \varphi^\alpha(t, \chi^\alpha), \quad \partial_\alpha \chi^\gamma|_{t=0} = \delta_\alpha^\gamma. \quad (50)$$

Then

$$\Omega_{\alpha\beta}(t, x) = \partial_\alpha \chi^\gamma \Omega_{\gamma\delta}^{(0)}(\chi) \partial_\beta \chi^\delta, \quad (51)$$

where the matrix $\Omega_{\alpha\beta}^{(0)}$ is the initial condition for $\Omega_{\alpha\beta}$,

$$\Omega_{\alpha\beta}(t, x)|_{t=0} = \Omega_{\alpha\beta}^{(0)}(x).$$

One can see that choosing the matrix $\Omega_{\alpha\beta}^{(0)}(x)$ to be nonsingular and subject to Jacobi identity, we guarantee the fulfilment of the same conditions for the complete matrix $\Omega_{\alpha\beta}(t, x)$ since components of the latter are given by a change of variables (51).

Thus, we see that for any set of first order equations (39), an integrating multiplier always exists, i.e. there always exist a Lagrangian $L(t, x, \dot{x})$ which has the set of equations

$$\Omega_{\alpha\beta}(\dot{x}^\beta - f^\beta(t, x)) = 0 \quad (52)$$

as the Euler-Lagrange equations. It is easy to see that this Lagrangian is linear in the first time derivative \dot{x}^α since equations (52) do not contain second-time derivatives, i.e. corresponding term $\partial^2 L / \partial \dot{x}^\alpha \partial \dot{x}^\beta$ vanishes. The general form of this Lagrangian is

$$L = J_\alpha \dot{x}^\alpha - H, \quad (53)$$

where $J_\alpha = J_\alpha(t, x)$ and $H = H(t, x)$ are some functions of the indicated arguments. The Euler-Lagrange equations corresponding to (53) are

$$\frac{\delta S}{\delta x} = \frac{\partial L}{\partial x} - \frac{d}{dt} \frac{\partial L}{\partial \dot{x}} = 0 \implies -\partial_\alpha H - \partial_t J_\alpha + (\partial_\alpha J_\beta - \partial_\beta J_\alpha) \dot{x}^\beta = 0. \quad (54)$$

Comparing equations (52) and (54) one gets²

$$\Omega_{\alpha\beta} = \partial_\alpha J_\beta - \partial_\beta J_\alpha, \quad (55)$$

and

$$\Omega_{\alpha\beta} f^\beta - \partial_t J_\alpha = \partial_\alpha H. \quad (56)$$

The functions J_α and H can be found from conditions (55) and (56) if the matrix $\Omega_{\alpha\beta}$ is given. One can see that consistency conditions for these equations exactly give us equations (46)-(48) for an integrating multiplier $\Omega_{\alpha\beta}$. We remind that the general solution for $J_\alpha(t, x)$ of the equation (55) provided that $\Omega_{\alpha\beta}$ is a given antisymmetric matrix that obeys the Jacobi identity, is given by

$$J_\alpha(t, x) = \int_0^1 x^\beta \Omega_{\beta\alpha}(t, sx) ds + \partial_\alpha \varphi(t, x), \quad (57)$$

where $\varphi(x)$ is an arbitrary function. Substituting (51) into (57), we obtain

$$J_\alpha(t, y) = \int_0^1 y^\beta \left[\partial_\alpha \chi^\gamma \Omega_{\gamma\delta}^{(0)}(\chi) \partial_\beta \chi^\delta \right]_{x=sy} ds + \partial_\alpha \varphi(t, y). \quad (58)$$

Equation (58) describes all the ambiguity (arbitrary symplectic matrix $\Omega_{\gamma\delta}^{(0)}$ and arbitrary function $\varphi(t, x)$) in constructing the term $J_\alpha(t, x)$ of the Lagrange function (53).

To restore the term H in the Lagrange function (53), we need to solve the equation (56) with respect to H . To this end, we remind that the general solution of the equation $\partial_i f = g_i$, provided a vector g_i is a gradient, is given by

$$f(x) = \int_0^1 ds x^i g_i(sx) + c,$$

where c is a constant. Taking the above into account, we obtain for H the following representation:

$$H(t, x) = \int_0^1 ds x^\beta [\Omega_{\beta\alpha}(t, sx) f^\alpha(t, sx) - \partial_t J_\beta(t, sx)] + c(t), \quad (59)$$

where $c(t)$ is an arbitrary function of time, and $\Omega_{\beta\alpha}$ and J_β are given by (51) and (58) respectively. All the arbitrariness in constructing H is thus due to arbitrary symplectic matrix $\Omega_{\gamma\delta}^{(0)}$, arbitrary functions $\varphi(t, x)$ entering into $\Omega_{\beta\alpha}$ and J_β and due $c(t)$.

²As a more abstract argument in favour of such a form of Ω we can give a Poincare Lemma, according to which any closed form is locally exact, but here we do not treat the global problems which can arise from the non-trivial topology of x^α -space.

We can see that there exist a family of Lagrangians (53) which lead to the same equations of motion (39). It is easy to see that actions with the same $\Omega_{\gamma\delta}^{(0)}$ but different functions $\varphi(t, x)$ and $c(t)$ differs by a total time derivative (we call such a difference trivial). A difference in Lagrange functions related to different choice of symplectic matrices $\Omega_{\alpha\beta}^{(0)}$ is not trivial. The corresponding Lagrangians are known as s -equivalent Lagrangians.

As an example, let us consider a theory with equations of motion of the form³

$$\dot{x} = A(t)x + j(t). \quad (60)$$

An action principle for such a theory can be constructed (see [13]) following the above described manner.

The solution of the Cauchy problem for the equations (60) reads

$$x(t) = \Gamma(t)x_{(0)} + \gamma(t), \quad (61)$$

where the matrix $\Gamma(t)$ is the fundamental solution of (60), i.e.,

$$\dot{\Gamma} = A\Gamma, \quad \Gamma(0) = 1, \quad (62)$$

and $\gamma(t)$ is a partial solution of (60). Then following (51), we construct the matrix Ω^4 ,

$$\Omega = \Lambda^T \Omega^{(0)} \Lambda, \quad \Lambda = \Gamma^{-1}. \quad (63)$$

and find the functions J and H according to (58) and (59),

$$J = \frac{1}{2}x\Omega, \quad H = \frac{1}{2}xBx - Cx, \quad (64)$$

where

$$B = \frac{1}{2}(\Omega A - A^T \Omega), \quad C = \Omega j. \quad (65)$$

Thus, the action functional for the general quadratic theory is

$$S[x] = \frac{1}{2} \int dt (x\Omega\dot{x} - xBx - 2Cx). \quad (66)$$

In conclusion we note that it is always possible to construct a Lagrangian action for any set of non-Lagrangian equations in an extended configuration space following a simple idea first proposed by Bateman [20]. Such a Lagrangian has the form of a sum of initial equations of motion being multiplied by the corresponding Lagrangian multipliers, new variables. The Euler-Lagrange equations for such an action contain besides the initial equations some new equations of motion for the Lagrange multipliers. In such an approach one has to think how to interpret the new variables already on the classical level. Additional difficulties (indefinite metric) can appear in course of the quantization [21]-[24].

Acknowledgement 1 *Gitman is grateful to the Brazilian foundations FAPESP and CNPq for permanent support; Kupriyanov thanks FAPESP for support.*

References

- [1] H. von Helmholtz, Journ. f. d. reine u. angew. Math., **100** (1887) 137.
- [2] G. Darboux, Lesons sur la Theorie Generale des Surfaces (Gauthier-Villars, Paris, 1894).
- [3] J. Douglas, Trans. Am. Math. Soc.50, N 71 (1941) 71.
- [4] P. Havas, Actra. Phys. Aust. **38** (1973) 145.
- [5] R. Santilli, Ann. Phys. (NY)**103** (1977) p.354.

³Here we use matrix notation, $x = (x^\alpha)$, $A(t) = (A(t)^\alpha_\beta)$, $j(t) = (j(t)^\alpha)$, $\alpha, \beta = 1, \dots, 2n$.

⁴For simplicity we choose the matrix $\Omega^{(0)}$ to be the constant nonsingular antisymmetric matrix.

- [6] W. Sarlet, *J.Math.Phys.***19**(1978)1049.
- [7] V.V. Dodonov, V.I. Man'ko and V.D. Skarzhinsky, *Arbitrariness in the choice of action and quantization of the given classical equations of motion*, preprint of P.N. Lebedev Physical Institute (1978).
- [8] S.Okubo, *Phys.Rev.* **D22**(1980)919.
- [9] W. Sarlet, *J.Phys.A* **15** (1982) 1503.
- [10] M. Henneaux, *Ann. Phys.* **140** (1982) 45.
- [11] G. Morandi, C. Ferrario, G. Lo Vecchio, G. Marmo and C. Rubano, *Phys. Rep.* **188** (1990) 147.
- [12] V.G. Kupriyanov, *Int.J.Theor.Phys.***45** (2006) 1129.
- [13] D.M. Gitman, V.G. Kupriyanov, *Eur.Phys.J.C***50**:691-700,2007.
- [14] A.P.Balachandran, T.R.Govindrajan and B.Vijayalakshimi, *Phys.Rev.* **D18**(1978)1950.
- [15] S. Hojman and L. Urrutia, *J. Math. Phys.* **22** (1981) 1896.
- [16] S.Hojman and H.Handerston, *J.Math.Phys.* **22** (1981) 1414.
- [17] M.Henneaux, L.Shepplley, *J.Math.Phys* **23** (1982) 2101.
- [18] J. Cislo and J. Lopuzanski, *J.Math.Phys.* **42** (2001) 5163.
- [19] P. Tempesta, E. Alfinito, R.Leo and G.Soliani, *J.Math.Phys.* **43** (2002) 3583.
- [20] H. Bateman, *Phys.Rev.* **38** (1931) 815.
- [21] H. Dekker, *Phys.Rep.* **80** (1981) 1.
- [22] E. Celeghini, M. Rasetti and G. Vitello, *Ann.Phys.* **215** (1992) 156.
- [23] R. Banerjee and P. Mukherjee, *J.Phys.A***35** (2002) 5591.
- [24] M. Blasone and P. Jizba, *Ann.Phys.* **312** (2004) 354.